

A NOISY SYSTEM WITH A FLATTENED HAMILTONIAN AND MULTIPLE TIME SCALES

NATELLA V. O'BRYANT

*Department of Mathematics, University of California - Irvine
272 Multipurpose Science & Technology Building, Irvine, CA 92697*

We consider a two-dimensional weakly dissipative dynamical system with time-periodic drift and diffusion coefficients. The average of the drift is governed by a degenerate Hamiltonian whose set of critical points has an interior. The dynamics of the system is studied in the presence of three time scales. Using the martingale problem approach and separating the time scales, we average the system to show convergence to a Markov process on a stratified space. The averaging combines the deterministic time averaging of periodic coefficients, and the stochastic averaging of the resulting system. The corresponding strata of the reduced space are a two-sphere, a point and a line segment. Special attention is given to the description of the domain of the limiting generator, including the analysis of the gluing conditions at the point where the strata meet. These gluing conditions, resulting from the effects of the hierarchy of time scales, are similar to the conditions on the domain of skew Brownian motion and are related to the description of spider martingales.

Keywords: Hamiltonian systems, Markov processes, stochastic averaging, martingale problem.

1. Introduction

Scientists have been studying Hamiltonian dynamical systems for centuries. One of the flourishing directions emerging as a result of these studies is current research on problems involving oscillations and vibrations of real mechanical systems. Generally, models that reflect such processes of real life are nonlinear and stochastic, and they often require some model reduction. One specific technique of model reduction is stochastic averaging which reduces more complicated random processes to simpler ones. We are interested in studying both the means of approximation of the initial model by its reduced model and the nature of the reduced model. One of the features of the reduced model is the structure of its state space, which may acquire dimensional discontinuities becoming a stratified space [?]. In cases like these, the domains of the generator of the limiting reduced process may require gluing conditions. The dynamics of the reduced process can be understood via the Fokker-Planck equation, with the gluing conditions reflected in the corresponding conservation of flux and continuity equations.

Real-life dynamical systems are typically described with equations which have small parameters. As these parameters vary, they can cause radical changes in the

qualitative structure of solutions. Although invisible on a usual time scale, in many cases these changes may become dramatically visible on a larger time scale. Bogolyubov [?] proved a deterministic result for ordinary differential equations which resolves a difficulty of this type, and which is known as the averaging principle.

The fact that most physical problems are non-deterministic inspired the development of stochastic averaging. The first rigorous arguments for stochastic averaging were given by Has'minskii [?], [?]. The essence of the classical stochastic averaging is the asymptotic separation of time scales.

Consider the following diffusion on $\mathbb{R}^2$:

$$\begin{aligned}\dot{\mathbf{X}}_t^\varepsilon(\mathbf{x}) &= \nabla^\perp H(\mathbf{X}_t^\varepsilon(\mathbf{x})) + \varepsilon \dot{\mathbf{W}}_t \\ \mathbf{X}_0^\varepsilon(\mathbf{x}) &= \mathbf{x},\end{aligned}$$

where $\nabla^\perp$ denotes a vector orthogonal to ∇ , H is a sufficiently smooth function, ε is a small parameter, and $\mathbf{W}$ is two-dimensional Brownian motion. Here H is the first integral of the corresponding deterministic system

$$\begin{aligned}\dot{\mathbf{X}}_t^0(\mathbf{x}) &= \nabla^\perp H(\mathbf{X}_t^0(\mathbf{x})) \\ \mathbf{X}_0^0(\mathbf{x}) &= \mathbf{x},\end{aligned}$$

This system is Hamiltonian with one degree of freedom. If the energy function H is a single-well Hamiltonian, i.e., if all the level sets $L_h = \{\mathbf{x} \in \mathbb{R}^2 : H(\mathbf{x}) = h\}$ of H have a single connected component formed by periodic trajectories of the system, then the invariant measure of this system depends only on the height h . There are two time scales involved here. The movement of the deterministic system is fast, but the energy of the stochastic system is varying slowly, and this slow transversal movement is visible only on a larger time scale. It is natural to separate the two time scales in order to witness the energy dynamics. From [?], on a large time scale, $H(\mathbf{X}^\varepsilon)$ weakly converges to a Markov process on a reduced space, which is the real line. In this case, the stratified space has one one-dimensional stratum.

Freidlin and Wentzell [?], and then Freidlin and Weber [?], considered the problem with a multiple-well Hamiltonian H . One of the essential assumptions made on the system in their work was the non-degenerate structure of the Hamiltonian. In the case of a multiple-well energy function, the invariant measure of the system depends not only on the height, but also on the particular component of the Hamiltonian. In the presence of saddle points the reduced space becomes a tree. Each vertex of the tree corresponds either to a single critical point of H , or to a whole homoclinic trajectory of the system. Every interior point of a segment on the tree corresponds to a particular periodic orbit. The energy of the stochastic system is varying slowly, and this slow transversal movement is naturally visible on a large time scale. Studying the energy dynamics as $\varepsilon \rightarrow 0$, the authors identified the limiting graph-valued process. They showed that the classical calculations of Has'minskii [?] can be applied, as long as the process stays away from the vertices of the graph. As for the vertices, the limiting Markov process requires gluing conditions to be enforced on the domain of its generator. The dynamics of the limiting process can be understood via the Fokker-Plank equation. The gluing reflects the

two following properties. The solution of the Fokker-Plank equation for the corresponding density must be continuous at vertices, and the weighted fluxes through vertices from different legs of a tree must sum to zero.

One way to describe a diffusion is to illustrate the dynamics of its trajectories by a corresponding stochastic differential equation. Another way, introduced by Stroock and Varadhan [?], is to build a martingale on the corresponding coordinate space. This alternative method is used to construct a limiting process by constructing a martingale problem for which this process is a solution. This is known as the martingale problem approach.

Sowers [?] investigated a stochastic system with a flattened Hamiltonian, removing the assumption of non-degeneracy. A canonical example of a Hamiltonian of this kind is

$$H(\mathbf{x}) = \left(\max \{0, \|\mathbf{x}\| - 1\} \right)^n, \quad n > 2.$$

To regularize noise, the chain equivalence [?] was used. This general equivalence relation identifies points if they can be taken from one to the other and back by a combination of small diffusive perturbation and fast drift. In this problem the limiting space has a dimensional discontinuity. The author showed convergence to a Markov process on the reduced space, whose strata are a two-sphere, a line segment and a point. A step towards a general theory of Markov processes on stratified spaces was made in the recent work of Evans and Sowers [?].

Recently, Namachchivaya and Sowers [?] made an attempt to develop a unified approach to study the energy dynamics of a non-degenerate single-degree-of-freedom system excited by both periodic and random perturbations. As a prototype for one of the cases, they studied a two-dimensional weakly dissipative system with time-periodic coefficients, and achieved model reduction through stochastic averaging.

We combine [?] and [?]. We consider a two-dimensional weakly dissipative system with time-periodic coefficients whose time average is governed by a degenerate Hamiltonian whose set of critical points has an interior. The dynamics of the system is studied in the presence of three time scales. The periodic fluctuations of the coefficients occur on the time scale of order $1/\varepsilon^2$, the effect of the drift is visible on the time scale of order $1/\varepsilon$, and the diffusion coefficients are of order one.

The main technical difficulty in this problem is stochastic averaging near the boundary of the critical set. The main step of the averaging is to approximately solve a certain partial differential equation. This equation involves drift of order one and small diffusion leading to a singular perturbation problem. We solve this problem by introducing new coordinates in the vicinity of the boundary. This coordinate transformation was inspired by Has'minskiĭ [?], and was previously used in [?]. These coordinates were introduced with the idea that equal angular displacement should correspond to equal amounts of diffusion across the boundary of the critical set. They perform a role similar to action-angle components. Solving the partial differential equation in Has'minskiĭ's coordinates involves Fourier analysis and leads to an inhomogeneous Bessel's ordinary differential equation. The unique solution is expressed via Bessel functions of small order and purely imaginary argument.

We perform stochastic averaging using the martingale problem. This approach was developed by Papanicolaou and Kohler [?]. Using the martingale problem and

separating the time scales, we average the system to show convergence to a Markov process on a stratified space. The corresponding strata are, as in [?], a two-sphere, a point and a line segment. Special attention is given to the description of the domain of the limiting generator, including the analysis of the gluing condition at the point where the strata meet. The differential operator governing the limiting process and the gluing describing the domain of its generator are given explicitly (Section 4.1). The gluing condition, resulting from the effects of the hierarchy of time scales, is similar to the conditions described in [?]. It is also similar to the conditions on the domain of skew Brownian motion, and is related to the description of spider martingales [?].

In our problem, the particular interest is in a careful analysis of the gluing condition on the domain of the limiting generator. Here, the multiple time scales affect gluing in a straightforward way. It would be interesting to see the changes to the initial problem which result in various changes to the structure of the reduced stratified space and which affect gluing in a different way.

1.1. Notation

Let $\alpha = (\alpha_1, \alpha_2)$ be an ordered tuple of nonnegative integers, i.e., let α be a multi-index. We set $|\alpha| = \alpha_1 + \alpha_2$ and

$$\partial^\alpha = \left(\frac{\partial}{\partial x_1} \right)^{\alpha_1} \left(\frac{\partial}{\partial x_2} \right)^{\alpha_2}.$$

We denote by $C^k(A; B)$ the collection of all functions $f : A \rightarrow B$ which have continuous derivatives of orders $0, 1, \dots, k$. When B is clear from the context, we write simply $C^k(A)$. We say that a function f is a C^k -diffeomorphism if f is a C^k -homeomorphism from A to B with a C^k inverse. We say that $f(\mathbf{x}, t) : A \times T \rightarrow B$ is $C^{k,l}(A \times T; B)$ if $\partial^\alpha f$ is continuous for all $|\alpha| \leq k$ and $\left(\frac{\partial f}{\partial t} \right)^\beta$ is continuous for all $0 \leq \beta \leq l$. If $l = 0$ we write $C^k(A \times T; B)$, and if $l = k = 0$ we write simply $C(A \times T; B)$. We denote by $C_c^k(A)$ the subset of $C^k(A)$ of functions having compact support, and by $C_b(A)$ the subset of continuous functions $C(A)$ bounded on the set A .

Let U be an open subset of $\mathbb{R}^2$. We denote by $\mathbb{H}^k(U)$ the Sobolev space, which consist of all locally integrable functions $h : U \rightarrow \mathbb{R}$ such that for each multi-index α with $|\alpha| \leq k$, $\partial^\alpha f$ exists in a weak sense and belongs to the space of square-integrable functions $L^2(U)$. For every $h \in \mathbb{H}^k(U)$, we define

$$\|h\|_{\mathbb{H}^k(U)} := \left(\sum_{|\alpha| \leq k} \int_U |\partial^\alpha h|^2 \right)^{1/2}.$$

For any functions $f \in C^{k,l}(\mathbb{R}^2 \times \mathbb{R}; \mathbb{R})$ and $g \in C^k(\mathbb{R}; \mathbb{R})$ ($k, l \geq 0$) and for any compact subsets $A \subset \mathbb{R}^2$, and $B, T \subset \mathbb{R}$, we define

$$\|f\|_{C^k(A \times T)} := \sup_{\substack{\mathbf{x} \in A \\ t \in T \\ |\alpha| \leq k}} |\partial^\alpha f(\mathbf{x}, t)|$$

and

$$\|g\|_{C^k(B)} := \sup_{\substack{\mathbf{y} \in B \\ |\alpha| \leq k}} |\partial^\alpha g(\mathbf{y})|.$$

We say that a function $f \in C^2(\mathbb{R}^2)$ is non-degenerate at a point $\mathbf{x}$ if the Hessian

$$D^2 f(\mathbf{x}) = \begin{vmatrix} \frac{\partial^2 f}{\partial x_1 \partial x_1}(\mathbf{x}) & \frac{\partial^2 f}{\partial x_1 \partial x_2}(\mathbf{x}) \\ \frac{\partial^2 f}{\partial x_2 \partial x_1}(\mathbf{x}) & \frac{\partial^2 f}{\partial x_2 \partial x_2}(\mathbf{x}) \end{vmatrix}$$

of f at $\mathbf{x}$ is non-degenerate, that is, if $\text{rank } D^2 f(\mathbf{x}) = 2$. We call a function $f \in C^2(\mathbb{R}^2)$ degenerate at $\mathbf{x}$ if $\text{rank } D^2 f(\mathbf{x}) < 2$.

2. Statement of the Problem

Since we will deal with a perturbed dynamical system, it makes sense to begin by describing our problem from the point of view of the theory of dynamical systems. To show that a certain limiting system is Markovian, we use the martingale problem, and we will need to restate our problem in the language more appropriate for this approach.

2.1. Mechanical description of the problem

When we speak about small perturbations, we will mean that the perturbing effect depends on a small parameter ε . Consider small random perturbations of a two-dimensional weekly-dissipative system with time-periodic coefficients. In other words, our interest is a system which has both periodic and stochastic perturbations,

$$\begin{aligned} d\tilde{\mathbf{X}}_t^\varepsilon &= \varepsilon b(\tilde{\mathbf{X}}_t^\varepsilon, t) dt + \varepsilon \sigma(\tilde{\mathbf{X}}_t^\varepsilon, t) d\mathbf{W}_t, \quad t \geq 0 \\ \tilde{\mathbf{X}}_0^\varepsilon &= \mathbf{x} \text{ a.s.} \end{aligned} \tag{2.1}$$

Here $\mathbf{W}$ is a two-dimensional Brownian motion. Coefficients b and σ in (2.1) are $C^{2,1}(\mathbb{R}^2 \times \mathbb{R})$ -functions and are 2π -periodic in time. We also assume that, given any $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2$,

$$(\mathbb{M}b)(\mathbf{x}) := \frac{1}{2\pi} \int_0^{2\pi} b(\mathbf{x}, t) dt = \nabla^\perp H(\mathbf{x}), \tag{2.2}$$

where $\nabla^\perp H(\mathbf{x}) = \left(\frac{\partial H(\mathbf{x})}{\partial x_2}, -\frac{\partial H(\mathbf{x})}{\partial x_1} \right)$ is perpendicular to $\nabla H(\mathbf{x})$. We assume that $H \geq 0$ is smooth and that it has a single well, that is, its level sets have only one connected component. Typically one assumes that H is non-degenerate at its critical points. Here we allow H to be degenerate at its critical points, so that $\text{rank } D^2 H(\mathbf{x}) \leq 2$, and that the set of the critical points of H can have an interior.

We will refer to the quantity H as the energy of the system (2.1). To describe H with some rigor, we introduce a function $K \in C^\infty(\mathbb{R}^2; \mathbb{R})$ with the following properties:

- (i) The set $\{\mathbf{x} \in \mathbb{R}^2 : K(\mathbf{x}) \leq 0\}$ is a diffeomorphism of the unit disc.

N. V. O'Bryant

- (ii) K is nonsingular on the set $\{\mathbf{x} \in \mathbb{R}^2 : K(\mathbf{x}) \geq 0\}$, meaning that $\nabla K(\mathbf{x}) \neq \mathbf{0}$ on this set.
- (iii) $\lim_{\|\mathbf{x}\| \rightarrow \infty} K(\mathbf{x}) = \infty$.

Having in mind these properties of K , we set

$$H(\mathbf{x}) := \left(\max \{0, K(\mathbf{x})\} \right)^n, n > 2.$$

We denote

$$\mathbf{V} := \{\mathbf{x} \in \mathbb{R}^2 : K(\mathbf{x}) < 0\}.$$

Example 2.1. Let K satisfy the above properties, $K(\mathbf{x}) = \|\mathbf{x}\| - 1$ for $\|\mathbf{x}\| \geq 1$ and $K(\mathbf{x}) < 0$ for $\|\mathbf{x}\| < 1$. Then $H(\mathbf{x}) = \left(\max \{0, \|\mathbf{x}\| - 1\} \right)^n$, $n > 2$, does not have any isolated critical points (Fig. 1). Instead, the set $\overline{\mathbf{V}}$ of its critical points




Fig. 1. H : View from below.

has an interior $\mathbf{V} = \{\mathbf{x} \in \mathbb{R}^2 : K(\mathbf{x}) < 0\} = \{\mathbf{x} \in \mathbb{R}^2 : \|\mathbf{x}\| < 1\}$. For any $\mathbf{x} \in \overline{\mathbf{V}}$, $\text{rank } D^2 H(\mathbf{x}) = 0$, so all the critical points of H are degenerate. Any positive level set $L_h := \{\mathbf{x} \in \mathbb{R}^2 : H(\mathbf{x}) = h\}$, $h > 0$, of H is a connected closed smooth curve in $\mathbb{R}^2$.

Remark 2.1. The property (ii) is essential for our construction. Set

$$\overline{K}(\mathbf{x}) = \begin{cases} 0 & \text{if } \|\mathbf{x}\| \leq 1, \\ \|\mathbf{x}\| \exp \left\{ \frac{1}{1-\|\mathbf{x}\|^2} \right\} & \text{if } \|\mathbf{x}\| > 1. \end{cases}$$

This smooth function satisfies the properties (i) and (iii), but not (ii), since it is singular on the set $\{\mathbf{x} \in \mathbb{R}^2 : \overline{K}(\mathbf{x}) = 0\}$.

The fact that K is nonsingular on $\{\mathbf{x} \in \mathbb{R}^2 : K(\mathbf{x}) \geq 0\}$ implies that there exists $a > 0$ such that the small set

$$\mathcal{E} = \{\mathbf{y} \in \mathbb{R}^2 : |K(\mathbf{y})| < a\} \tag{2.3}$$

surrounding the boundary $\partial\mathbf{V}$ of the set $\overline{\mathbf{V}} = \{\mathbf{x} \in \mathbb{R}^2 : K(\mathbf{x}) \leq 0\}$ has one component and $\nabla K(\mathbf{x}) \neq \mathbf{0}$ for all $\mathbf{x} \in \overline{\mathcal{E}}$.

Our goal is to study the dynamics of the energy of the system (2.1) as the small parameter ε tends to 0 and to describe the limiting system. We will show that it has the Markovian property.

Definition 2.1. We write $\gamma = \sigma\sigma^T$, and for any functions $f, g \in C^2(\mathbb{R}^2; \mathbb{R})$ and for every $t \in \mathbb{R}$, we define two differential operators: the generator

$$(\mathcal{L}_t f)(\mathbf{x}) := \frac{1}{2} \sum_{i,j=1}^2 \gamma_{ij}(\mathbf{x}, t) \frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{x}); \quad (2.4)$$

and the bracket

$$\langle df, dg \rangle_t(\mathbf{x}) := \sum_{i,j=1}^2 \gamma_{ij}(\mathbf{x}, t) \frac{\partial f}{\partial x_i}(\mathbf{x}) \frac{\partial g}{\partial x_j}(\mathbf{x}). \quad (2.5)$$

If $f, g \in C^{2,1}(\mathbb{R}^2 \times \mathbb{R}; \mathbb{R})$, then $(\mathcal{L}_t f)(\mathbf{x}, t)$ and $\langle df, dg \rangle_t(\mathbf{x}, t)$ are obtained by applying $\mathcal{L}_t$ and $\langle df, dg \rangle_t$ to $f(\cdot, t)$ and $g(\cdot, t)$, i.e., $(\mathcal{L}_t f)(\mathbf{x}, t) = (\mathcal{L}_t f(\cdot, t))(\mathbf{x})$ and $\langle df, dg \rangle_t f(\mathbf{x}, t) = (\langle df, dg \rangle_t(\cdot, t))(\mathbf{x})$.

Remark 2.2. If $h \in C^2(\mathbb{R}; \mathbb{R})$ and $f \in C^2(\mathbb{R}^2; \mathbb{R})$ then

$$\left(\mathcal{L}_t (h \circ f) \right)(\mathbf{x}) = h'(f(\mathbf{x}))(\mathcal{L}_t f)(\mathbf{x}) + \frac{1}{2} h''(f(\mathbf{x})) \langle df, df \rangle_t(\mathbf{x}).$$

Applying Itô's formula, from (2.1) we have that

$$\begin{aligned} H(\tilde{\mathbf{X}}_t^\varepsilon) &= H(\mathbf{x}) + \varepsilon \int_0^t \left(\nabla H(\tilde{\mathbf{X}}_s^\varepsilon), b(\tilde{\mathbf{X}}_s^\varepsilon, s) \right) ds \\ &\quad + \varepsilon \int_0^t \left(\nabla H(\tilde{\mathbf{X}}_s^\varepsilon), \sigma(\tilde{\mathbf{X}}_s^\varepsilon, s) \right) d\mathbf{W}_s + \varepsilon^2 \int_0^t (\mathcal{L}_s H)(\tilde{\mathbf{X}}_s^\varepsilon) ds, \end{aligned}$$

and we see that $H(\tilde{\mathbf{X}}^\varepsilon)$ is slowly varying (as opposed to being conserved). We want to choose a time scale fine enough to see the effect of the periodic coefficient b on the fluctuations of $H(\tilde{\mathbf{X}}^\varepsilon)$. In order to do this, we rescale time by introducing the process $\mathbf{X}^\varepsilon$ so that $\mathbf{X}_t^\varepsilon = \tilde{\mathbf{X}}_{t/\varepsilon^2}^\varepsilon$ for every $t \geq 0$. Using the scaling property of Brownian motion, without loss of generality, we can rewrite (2.1) in the form

$$\begin{aligned} d\mathbf{X}_t^\varepsilon &= \frac{1}{\varepsilon} b \left(\mathbf{X}_t^\varepsilon, \frac{t}{\varepsilon^2} \right) dt + \sigma \left(\mathbf{X}_t^\varepsilon, \frac{t}{\varepsilon^2} \right) d\mathbf{W}_t, \quad t \geq 0 \\ \mathbf{X}_0^\varepsilon &= \mathbf{x} \text{ a.s.} \end{aligned} \quad (2.6)$$

We can distinguish between three time scales here. The motion described by (2.6) consists of the very fast (of order $1/\varepsilon^2$) periodic oscillation of the coefficients, the fast (of order $1/\varepsilon$) rotation along the trajectories of the corresponding unperturbed system, and the slow diffusion (of order 1) across these trajectories. From (2.6),

N. V. O'Bryant

using Itô's formula,

$$\begin{aligned} H(\mathbf{X}_t^\varepsilon) &= H(\mathbf{x}) + \frac{1}{\varepsilon} \int_0^t \left(\nabla H(\mathbf{X}_s^\varepsilon), b\left(\mathbf{X}_s^\varepsilon, \frac{s}{\varepsilon^2}\right) \right) ds \\ &\quad + \int_0^t \left(\nabla H(\mathbf{X}_s^\varepsilon), \sigma\left(\mathbf{X}_s^\varepsilon, \frac{s}{\varepsilon^2}\right) \right) d\mathbf{W}_s + \int_0^t (\mathcal{L}_{s/\varepsilon^2} H)(\mathbf{X}_s^\varepsilon) ds. \end{aligned}$$

After rescaling, we can rephrase our goal. We want to study the behavior of $H(\mathbf{X}^\varepsilon)$ as $\varepsilon \downarrow 0$, and to show that the process describing the energy of the limiting system is Markovian.

2.2. Topological identification and stratification of the quotient space

Let ξ_t be the flow generated by $\nabla^\perp H$, i.e.,

$$\dot{\xi}_t(\mathbf{x}) = \nabla^\perp H(\xi_t(\mathbf{x})), \quad \xi_0(\mathbf{x}) = \mathbf{x}.$$

We will use ξ_t to form a quotient space. Let $\sim$ be the chain equivalence relation based on ξ_t .

All trajectories of ξ_t are periodic on $\mathbb{R}^2 \setminus \overline{\mathbf{V}}$, therefore the chain orbit of $\mathbf{x} \in \mathbb{R}^2 \setminus \overline{\mathbf{V}}$ consists of all the points on the level set to which $\mathbf{x}$ belongs. All points of the open set $\mathbf{V}$ are fixed points of the flow ξ_t , and every point $\mathbf{x} \in \mathbf{V}$ is its own chain orbit. All points of $\partial\mathbf{V}$ are also fixed points of the flow, but the chain orbit of every such point is the whole set $\partial\mathbf{V}$, since every point of $\partial\mathbf{V}$ always has a neighborhood containing points from $\mathbb{R}^2 \setminus \overline{\mathbf{V}}$.

Fix a level of H of height $H^* > 0$, let $K^* := (H^*)^{1/n}$, and define the two open sets

$$\mathbf{U} := \{\mathbf{x} \in \mathbb{R}^2 : 0 < H(\mathbf{x}) < H^*\} = \{\mathbf{x} \in \mathbb{R}^2 : 0 < K(\mathbf{x}) < K^*\}$$

and

$$\mathbf{I} := \{\mathbf{x} \in \mathbb{R}^2 : H(\mathbf{x}) < H^*\} = \{\mathbf{x} \in \mathbb{R}^2 : K(\mathbf{x}) < K^*\}.$$

The sets $\mathbf{V}, \partial\mathbf{V}, \mathbf{U}, \partial\mathbf{I}$ are disjoint with union $\bar{\mathbf{I}}$ (Fig. 2).




Fig. 2. $\bar{\mathbf{I}} = \mathbf{V} \cup \partial\mathbf{V} \cup \mathbf{U} \cup \partial\mathbf{I}$.

Define

$$\Gamma := \bar{\mathbf{I}} / \sim.$$

Then Γ , endowed with the quotient topology defined by $\sim$, is a quotient space. For every $\mathbf{x} \in \bar{\mathbf{I}}$, define the natural map

$$\pi(\mathbf{x}) := [\mathbf{x}] = \{\mathbf{y} \in \bar{\mathbf{I}} : \mathbf{y} \sim \mathbf{x}\}.$$

Notice that

$$\pi(\mathbf{x}) = \begin{cases} H^{-1}(H(\mathbf{x})) & \text{if } \mathbf{x} \in \mathbf{U} \cup \partial\mathbf{I}, \\ \partial\mathbf{V} & \text{if } \mathbf{x} \in \partial\mathbf{V}, \\ \{\mathbf{x}\} & \text{if } \mathbf{x} \in \mathbf{V}. \end{cases}$$

For $A \subset \bar{\mathbf{I}}$ denote $\Gamma_A = \pi(A) = \{[\mathbf{x}] \in \Gamma : \mathbf{x} \in A\}$ so that $\Gamma = \Gamma_{\mathbf{V}} \cup \Gamma_{\partial\mathbf{V}} \cup \Gamma_{\mathbf{U}} \cup \Gamma_{\partial\mathbf{I}}$.

Proposition 2.1. *There is a homeomorphism from Γ to the submanifold*

$$\mathbb{S}^2 \cup \left(\{0\} \times \{0\} \times [1, H^* + 1] \right)$$

in $\mathbb{R}^3$, where $\mathbb{S}^2$ is the 2-sphere $\{\mathbf{x} \in \mathbb{R}^3 : \|\mathbf{x}\| = 1\}$.

Proof. Denote $\mathbb{S}^2 \cup \left(\{0\} \times \{0\} \times [1, H^* + 1] \right)$ by $\mathbf{M}_3$, and define the map $p : \Gamma \rightarrow \mathbf{M}_3$ by

$$p([\mathbf{x}]) := \begin{cases} s(\mathbf{x}) & \text{if } [\mathbf{x}] \in \Gamma_{\mathbf{V}}, \\ (0, 0, 1) & \text{if } [\mathbf{x}] \in \Gamma_{\partial\mathbf{V}}, \\ (0, 0, H(\mathbf{x}) + 1) & \text{if } [\mathbf{x}] \in \Gamma_{\mathbf{U}} \cup \Gamma_{\partial\mathbf{I}}, \end{cases}$$

where $s : \bar{\mathbf{V}} \rightarrow \mathbb{S}^2$ is a homeomorphism such that $s(\partial\mathbf{V}) = (0, 0, 1)$. Notice that p is the desired homeomorphism from Γ to $\mathbf{M}_3$. $\square$

We see that $\Gamma_{\mathbf{V}}$ is an open two-dimensional smooth manifold, $\Gamma_{\mathbf{U}}$ is an open one-dimensional smooth manifold, $\Gamma_{\mathbf{V}} \cap \Gamma_{\mathbf{U}} = \emptyset$. $\Gamma_{\partial\mathbf{V}}$ is a limit point of both $\Gamma_{\mathbf{V}}$ and $\Gamma_{\mathbf{U}}$, and $\Gamma_{\partial\mathbf{I}}$ is the other limit point of $\Gamma_{\mathbf{U}}$. Both $\Gamma_{\partial\mathbf{V}}$ and $\Gamma_{\partial\mathbf{I}}$ can be treated as zero-dimensional smooth manifolds, and the Whitney regularity condition is trivially satisfied. Therefore, $\Gamma = \Gamma_{\mathbf{V}} \cup \Gamma_{\partial\mathbf{V}} \cup \Gamma_{\mathbf{U}} \cup \Gamma_{\partial\mathbf{I}}$ is a stratified space.

2.3. Probabilistic formulation of the problem

Probabilistically, we are interested in the laws of a stochastic processes. We would like to show that a certain limiting law is Markovian in nature. The martingale problem is a standard tool for such analysis. We will restate our problem in the language natural for the martingale approach.

For any $f \in C^2(\mathbb{R}^2)$, and for every $t \in \mathbb{R}$, define the differential operator

$$(\mathcal{L}_t^\varepsilon f)(\mathbf{x}) := \frac{1}{\varepsilon} (\nabla f(\mathbf{x}), b(\mathbf{x}, t)) + (\mathcal{L}_t f)(\mathbf{x}),$$

with $\mathcal{L}_t f$ defined in (2.4) above. If $f \in C^{2,1}(\mathbb{R}^2 \times \mathbb{R})$, then $(\mathcal{L}_t^\varepsilon f)(\mathbf{x}, t)$ is obtained by applying $\mathcal{L}_t^\varepsilon$ to $f(\cdot, t)$ i.e., $(\mathcal{L}_t^\varepsilon f)(\mathbf{x}, t) = (\mathcal{L}_t f(\cdot, t))(\mathbf{x})$. We assume that the domain $\mathcal{D}^\varepsilon$ of the generator $\mathcal{L}_t^\varepsilon$ contains a dense subset of functions from $C^2(\mathbb{R}^2)$.

Let $\Omega := C([0, \infty); \mathbb{R}^2)$. Given $\mathbf{x} \in \bar{\mathbf{I}}$, let $\mathbb{P}^\varepsilon \in \mathcal{P}(C([0, \infty); \mathbb{R}^2))$ be a solution of the stopped martingale problem for $(\mathcal{L}_t^\varepsilon, \delta_{\mathbf{x}}, \mathbf{I}), t \geq 0$. Then for every $f \in C^{2,1}(\mathbb{R}^2 \times [0, \infty))$,

$$f(\mathbf{X}_{t \wedge \tau}, t \wedge \tau) - f(\mathbf{X}_{s \wedge \tau}, s \wedge \tau) - \int_{s \wedge \tau}^{t \wedge \tau} \left(\frac{\partial f}{\partial u}(\mathbf{X}_u, u) + (\mathcal{L}_u^\varepsilon f)(\mathbf{X}_u, u) \right) du$$

is a $\mathbb{P}^\varepsilon$ -martingale.

Let $\mathbf{X}$ be the corresponding coordinate process defined by $\mathbf{X}_t(\omega) := \omega(t)$, $t \geq 0$, $\omega \in \Omega$. Set $\mathcal{F}_t$ to be generated by this coordinate process up to time t , and define $\mathcal{F} := \bigvee_t \mathcal{F}_t$. Define the $\mathcal{F}_t$ -stopping time τ by

$$\tau := \inf \{s \geq 0 : \mathbf{X}_s \notin \mathbf{I}\} = \inf \{s \geq 0 : H(\mathbf{X}_s) \geq H^*\},$$

i.e., τ is defined to be the time of the first exit from $\mathbf{I}$.

Our condition on $\mathbb{P}^\varepsilon$ means that the corresponding coordinate process $\mathbf{X}$ on $(\Omega, \mathcal{F}, \mathbb{P}^\varepsilon)$ is a solution to the stopped martingale problem for $(\mathcal{L}_t^\varepsilon, \delta_{\mathbf{x}}, \mathbf{I}), t \geq 0$.

Define $\mathbf{Y}_t := [\mathbf{X}_{t \wedge \tau}]$, $t \geq 0$. For each $\varepsilon > 0$, define the probability measure $\mathbb{P}^{\varepsilon,*}(A) := \mathbb{P}^\varepsilon\{\mathbf{Y} \in A\}$, $A \in \mathcal{B}(C([0, \infty), \mathbf{I}))$. Let $\mathbf{Y}_t(\omega) = \omega(t)$, $t \geq 0$, $\omega \in C([0, \infty), \mathbf{I})$, be the corresponding coordinate process. For each $t \geq 0$, define $\mathcal{F}_t^* := \sigma\{\mathbf{Y}_s, 0 \leq s \leq t\}$, and define a σ -algebra on $\Omega^* = C([0, \infty), \mathbf{I})$ by $\mathcal{F}^* := \bigvee_{t \geq 0} \mathcal{F}_t^*$.

With this in mind, we can formulate our task once again; this time in the language for the martingale approach. Our goal is to show that the $\mathbb{P}^\varepsilon$ -laws of $\mathbf{Y}$ converge to the law of a $\mathbf{I}$ -valued Markov process. In other words, we want to understand the asymptotics of $\mathbb{P}^{\varepsilon,*}$ as $\varepsilon \downarrow 0$. Even more precisely, we will prove that $\mathbb{P}^{\varepsilon,*}$ converges to the unique solution $\mathbb{P}^*$ of the martingale problem for $(\mathcal{L}^*, \delta_{[\mathbf{x}]})$, and we will identify the limiting generator $\mathcal{L}^*$ and its domain.

3. Averaging

The main goal of our work is model reduction. To achieve this goal, we use both time averaging and the stochastic averaging of Freidlin and Wentzell as our main techniques.

We study our system in the presence of three time scales. The periodic fluctuations of the coefficients occur on the time scale of order $1/\varepsilon^2$, the effect of the drift is visible in time of order $1/\varepsilon$, and diffusion is of order one. The hierarchy of the time scales indicate that our system will require multiple-level averaging. First, we average the quickly-varying periodic coefficients.

3.1. First-level averaging

We recall that $\mathbb{P}^\varepsilon \in \mathcal{P}(C[0, \infty); \mathbb{R})$ is a solution of the stopped martingale problem for $(\mathcal{L}_t^\varepsilon, \delta_{\mathbf{x}}, \mathbf{I}), t \geq 0$, $\mathbf{x} \in \bar{\mathbf{I}}$, where $\mathbf{I} := \{\mathbf{x} \in \mathbb{R}^2 : H(\mathbf{x}) < H^*\}$, and $\tau := \inf \{s \geq 0 : \mathbf{X}_s \notin \mathbf{I}\}$. Let $\mathbb{E}^\varepsilon$ be the expectation operator associated with $\mathbb{P}^\varepsilon$.

Lemma 3.1. *If $f \in C^2(\bar{\mathbf{I}} \times [0, \infty); \mathbb{R})$ is 2π -periodic in its last argument, then there exists a positive constant C such that*

$$\mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left(f\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}f)(\mathbf{X}_u) \right) du \right| \right] \leq C\varepsilon(1+t) \|f\|_{C^2(\bar{\mathbf{I}} \times [0, \infty))}. \quad (3.7)$$

Proof. We define

$$\Phi_f(\mathbf{x}, t) := \int_0^t (f(\mathbf{x}, u) - (\mathbb{M}f)(\mathbf{x})) du. \quad (3.8)$$

For any $\mathbf{x} \in \bar{\mathbf{I}}$, any multi-index α and $t \geq 0$,

$$\begin{aligned} \partial^\alpha \Phi_f(\mathbf{x}, t) &= \int_0^t \left(\partial^\alpha f(\mathbf{x}, u) - \frac{1}{2\pi} \int_0^{2\pi} \partial^\alpha f(\mathbf{x}, s) ds \right) du \\ &= \int_{2\pi \lfloor \frac{t}{2\pi} \rfloor}^t \partial^\alpha f(\mathbf{x}, u) du - \left(\frac{t}{2\pi} - \lfloor \frac{t}{2\pi} \rfloor \right) \int_0^{2\pi} \partial^\alpha f(\mathbf{x}, u) du. \end{aligned}$$

This implies that for every $0 < k \leq 2$

$$\|\Phi_f\|_{C^k(\bar{\mathbf{I}} \times [0, \infty))} \leq 4\pi \|f\|_{C^k(\bar{\mathbf{I}} \times [0, \infty))}. \quad (3.9)$$

By the stopped martingale problem for $(\mathcal{L}_{t/\varepsilon^2}^\varepsilon, \delta_{\mathbf{x}}, \mathbf{I})$,

$$\begin{aligned} \Phi_f\left(\mathbf{X}_{t \wedge \tau}, \frac{t \wedge \tau}{\varepsilon^2}\right) - \Phi_f(\mathbf{x}, 0) &= \frac{1}{\varepsilon^2} \int_0^{t \wedge \tau} \frac{\partial \Phi_f}{\partial u}\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) du \\ &\quad + \frac{1}{\varepsilon} \int_0^{t \wedge \tau} \left(\nabla_{\mathbf{x}} \Phi_f\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right), b\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) \right) du \\ &\quad + \int_0^{t \wedge \tau} (\mathcal{L}_{u/\varepsilon^2} \Phi_f)\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) du \\ &\quad + \mathbf{M}_{t \wedge \tau}^{\Phi_f, \varepsilon}, \end{aligned} \quad (3.10)$$

where $\mathbf{M}^{\Phi_f, \varepsilon}$ is the $\mathbb{P}^\varepsilon$ -martingale with its quadratic variation

$$\langle \mathbf{M}^{\Phi_f, \varepsilon} \rangle_t = \int_0^t \langle d\Phi_f, d\Phi_f \rangle_{u/\varepsilon^2} \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) du.$$

Then

$$\begin{aligned} \int_0^{t \wedge \tau} \left(f\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}f)(\mathbf{X}_u) \right) du &= \varepsilon^2 \left(\Phi_f\left(\mathbf{X}_{t \wedge \tau}, \frac{t \wedge \tau}{\varepsilon^2}\right) - \Phi_f(\mathbf{x}, 0) \right) \\ &\quad - \varepsilon \int_0^{t \wedge \tau} \left(\nabla_{\mathbf{x}} \Phi_f\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right), b\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) \right) du \\ &\quad - \varepsilon^2 \int_0^{t \wedge \tau} (\mathcal{L}_{u/\varepsilon^2} \Phi_f)\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) du \\ &\quad - \varepsilon^2 \mathbf{M}_{t \wedge \tau}^{\Phi_f, \varepsilon}. \end{aligned} \quad (3.11)$$

By Burkholder-Davis-Gundy inequality, there exists a universal positive constant C_m such that

$$\mathbb{E}^\varepsilon \left[\left| \mathbf{M}_{t \wedge \tau}^{\Phi_f, \varepsilon} \right| \right] \leq C_m \mathbb{E}^\varepsilon \left[\langle \mathbf{M}^{\Phi_f, \varepsilon} \rangle_{t \wedge \tau}^{1/2} \right] \quad (3.12)$$

From (3.9), there exists a positive constant C such that

$$\langle \mathbf{M}^{\Phi_{f,\varepsilon}} \rangle_{t \wedge \tau} \leq Ct \|f\|_{C^1(\bar{\mathbf{I}} \times [0, \infty))}^2. \quad (3.13)$$

Finally, from expression (3.11), applying inequalities (3.9), (3.12), and (3.13), there is a positive constant C such that (3.7) holds. $\square$

3.2. Stochastic averaging away from the critical set

Let ϕ_t be the flow generated by $\nabla^\perp K$, i.e.,

$$\dot{\phi}_t(\mathbf{x}) = \nabla^\perp K(\phi_t(\mathbf{x})), \quad \phi_0(\mathbf{x}) = \mathbf{x}.$$

Here we are concerned with stochastic averaging on a subset of $\bar{\mathbf{I}} \setminus \bar{\mathbf{V}} = \mathbf{U} \cup \partial\mathbf{I}$. On any such subset, $\xi_t(\mathbf{x}) = \phi_{n(K(\mathbf{x}))^{n-1}t}(\mathbf{x})$ and the flow ϕ_t has the same orbits as the flow ξ_t generated by $\nabla^\perp H$. To average over these orbits, we define

$$\eta(K(\mathbf{x})) := \inf \{t > 0 : \phi_t(\mathbf{x}) = \mathbf{x}\},$$

and for every $f \in C(\bar{\mathbf{I}}; \mathbb{R})$ we define

$$\Psi_f(\mathbf{x}) := \frac{1}{\eta(K(\mathbf{x}))} \int_0^{\eta(K(\mathbf{x}))} t f(\phi_t(\mathbf{x})) dt. \quad (3.14)$$

Let $\mathcal{D}_{\mathbb{A}_0}$ be the set of all real-valued functions whose restriction to $\mathbf{V} \cup \mathbf{U}$ is continuous. We define the pre-averaging operator $\mathbb{A}_0 : \mathcal{D}_{\mathbb{A}_0} \rightarrow C(\Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}})$ by

$$(\mathbb{A}_0 f)([\mathbf{x}]) := \begin{cases} f(\mathbf{x}) & \text{if } [\mathbf{x}] \in \Gamma_{\mathbf{V}} \\ \left(\int_{\mathbf{y} \in [\mathbf{x}]} \frac{1}{\|\nabla K(\mathbf{y})\|} dl(\mathbf{y}) \right)^{-1} \int_{\mathbf{y} \in [\mathbf{x}]} \frac{f(\mathbf{y})}{\|\nabla K(\mathbf{y})\|} dl(\mathbf{y}) & \text{if } [\mathbf{x}] \in \Gamma_{\mathbf{U}}, \end{cases}$$

where dl is the arc-length element.

Let $\mathcal{D}_{\mathbb{A}}$ be the subset of functions f from $\mathcal{D}_{\mathbb{A}_0}$ for which

$$\lim_{\substack{[\mathbf{y}] \rightarrow \Gamma_{\partial\mathbf{V}} \\ [\mathbf{y}] \in \Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}}}} (\mathbb{A}_0 f)([\mathbf{y}]) \quad \text{and} \quad \lim_{\substack{[\mathbf{y}] \rightarrow \Gamma_{\partial\mathbf{U}} \\ [\mathbf{y}] \in \Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}}}} (\mathbb{A}_0 f)([\mathbf{y}])$$

exist. We define the averaging operator $\mathbb{A} : \mathcal{D}_{\mathbb{A}} \rightarrow C(\Gamma)$ by

$$(\mathbb{A} f)([\mathbf{x}]) := \lim_{\substack{[\mathbf{y}] \rightarrow [\mathbf{x}] \\ [\mathbf{y}] \in \Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}}}} (\mathbb{A}_0 f)([\mathbf{y}]).$$

Remark 3.3. We notice that if $\mathbf{x} \in \mathbf{U} \cup \partial\mathbf{I}$, then $[\mathbf{x}] = K^{-1}(K(\mathbf{x}))$, i.e., $[\mathbf{x}]$ is totally defined by $K(\mathbf{x})$, and $(\mathbb{A} f)([\mathbf{x}]) = (\mathbb{A} f)(K^{-1}(K(\mathbf{x})))$. We define $(\mathbb{A}_K f)(h)$ to be the average of f on the level set $\{\mathbf{x} \in \mathbb{R}^2 : K(\mathbf{x}) = h\}$, i.e.,

$$(\mathbb{A}_K f)(K(\mathbf{x})) := \frac{1}{\eta(K(\mathbf{x}))} \int_0^{\eta(K(\mathbf{x}))} f(\phi_t(\mathbf{x})) dt.$$

Then for any $\mathbf{x} \in \mathbf{U} \cup \partial\mathbf{I}$,

$$(\mathbb{A} f)([\mathbf{x}]) = (\mathbb{A}_K f)(K(\mathbf{x})).$$

Let the function $\omega \in C_c^\infty(\mathbb{R})$ be supported on a compact subset of $(0, \infty)$.

Lemma 3.2. *For any function $f \in C^3(\bar{\mathbf{I}}; \mathbb{R})$ and for any $0 < p < 1$ there exists a constant $C_a > 0$ such that*

$$\mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left(f(\mathbf{X}_u) - (\mathbb{A}f)([\mathbf{X}_u]) \right) \omega \left(\frac{K(\mathbf{X}_u)}{\varepsilon^p} \right) du \right| \right] < C_a \varepsilon^{1-p(n+2)} (1+t) \|f\|_{C^3(\bar{\mathbf{I}})}. \quad (3.15)$$

Proof. Let $\mathbf{x} \in \bar{\mathbf{I}}$. Since ω is supported on a compact subset of $(0, \infty)$, either

$$\left(f(\mathbf{x}) - (\mathbb{A}f)([\mathbf{x}]) \right) \omega \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) = 0,$$

or there exist some positive constants C_1 and C_2 such that $C_1 \varepsilon^p < K(\mathbf{x}) < C_2 \varepsilon^p$, and therefore we may assume that $\mathbf{x} \in \mathbf{U} \cup \partial \mathbf{I}$. From the above remark, this implies that $(\mathbb{A}f)([\mathbf{x}]) = (\mathbb{A}_K f)(K(\mathbf{x}))$.

From (3.14), there exist a constant $C > 0$ such that

$$\|\Psi_f\|_{C^k(\bar{\mathbf{I}})} < C \|f\|_{C^k(\bar{\mathbf{I}})}. \quad (3.16)$$

Notice that $\eta(K(\phi_t(\mathbf{x}))) = \eta(K(\mathbf{x}))$ and

$$\begin{aligned} \Psi_f(\phi_t(\mathbf{x})) &= \frac{1}{\eta(K(\mathbf{x}))} \int_0^{\eta(K(\mathbf{x}))} s f(\phi_s(\phi_t(\mathbf{x}))) ds \\ &= \frac{1}{\eta(K(\mathbf{x}))} \int_t^{t+\eta(K(\mathbf{x}))} (s-t) f(\phi_s(\mathbf{x})) ds. \end{aligned}$$

Differentiating the last equality, we obtain

$$\begin{aligned} \frac{d}{dt} \Psi_f(\phi_t(\mathbf{x})) &= \frac{1}{\eta(K(\mathbf{x}))} \frac{d}{dt} \left(\int_t^{t+\eta(K(\mathbf{x}))} (s-t) f(\phi_s(\mathbf{x})) ds \right) \\ &= f(\phi_t(\mathbf{x})) - (\mathbb{A}_K f)(K(\phi_t(\mathbf{x}))). \end{aligned} \quad (3.17)$$

On the other hand, using the fact that the flow ϕ_t is generated by $\nabla^\perp K$, we observe that

$$\frac{d}{dt} \Psi_f(\phi_t(\mathbf{x})) = (\nabla \Psi_f(\phi_t(\mathbf{x})), \nabla^\perp K(\phi_t(\mathbf{x}))), \quad (3.18)$$

Expressions (3.17) and (3.18) combined yield

$$f(\phi_t(\mathbf{x})) - (\mathbb{A}_K f)(K(\phi_t(\mathbf{x}))) = (\nabla \Psi_f(\phi_t(\mathbf{x})), \nabla^\perp K(\phi_t(\mathbf{x}))).$$

This equality, being true for any $t \in \mathbb{R}$, implies that for any $\mathbf{x} \in \bar{\mathbf{I}}$,

$$\left(f(\mathbf{x}) - (\mathbb{A}_K f)(K(\mathbf{x})) \right) \omega \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) = (\nabla \Psi_f(\mathbf{x}), \nabla^\perp K(\mathbf{x})) \omega \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right). \quad (3.19)$$

N. V. O'Bryant

For any $\mathbf{x} \in \bar{\mathbf{I}}$, we define

$$\Psi_f^\varepsilon(\mathbf{x}) := \frac{1}{n(K(\mathbf{x}))^{n-1}} \Psi_f(\mathbf{x}) \omega\left(\frac{K(\mathbf{x})}{\varepsilon^p}\right)$$

and notice that since $(\mathbb{M}b)(\mathbf{x}) = n(K(\mathbf{x}))^{n-1} \nabla^\perp K(\mathbf{x})$, then

$$\left(\mathbb{M}(\nabla \Psi_f^\varepsilon, b)\right)(\mathbf{x}) = (\nabla \Psi_f(\mathbf{x}), \nabla^\perp K(\mathbf{x})) \omega\left(\frac{K(\mathbf{x})}{\varepsilon^p}\right).$$

Using inequality (3.16), we see that there exists a constant $C > 0$, such that

$$\|\Psi_f^\varepsilon\|_{C^k(\bar{\mathbf{I}})} < C \varepsilon^{-p(n-1+k)} \|f\|_{C^k(\bar{\mathbf{I}})}. \quad (3.20)$$

This implies that there exists a positive constant C such that

$$\|(\nabla \Psi_f^\varepsilon, b)\|_{C^2(\bar{\mathbf{I}} \times [0, \infty))} < C \varepsilon^{-p(n+2)} \|f\|_{C^3(\bar{\mathbf{I}})}. \quad (3.21)$$

By the stopped martingale problem for $(\mathcal{L}_{t/\varepsilon^2}^\varepsilon, \delta_{\mathbf{x}}, \mathbf{I})$,

$$\begin{aligned} \int_0^{t \wedge \tau} \left(\nabla \Psi_f^\varepsilon(\mathbf{X}_u), b\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) \right) du &= \varepsilon (\Psi_f^\varepsilon(\mathbf{X}_{t \wedge \tau}) - \Psi_f^\varepsilon(\mathbf{x})) \\ &\quad - \varepsilon \int_0^{t \wedge \tau} (\mathcal{L}_{u/\varepsilon^2}^\varepsilon \Psi_f^\varepsilon)(\mathbf{X}_u) du - \varepsilon \mathbf{M}_{t \wedge \tau}^{\Psi_f^\varepsilon, \varepsilon}, \end{aligned} \quad (3.22)$$

where $\mathbf{M}^{\Psi_f^\varepsilon, \varepsilon}$ is the $\mathbb{P}^\varepsilon$ -martingale with its quadratic variation

$$\langle \mathbf{M}^{\Psi_f^\varepsilon, \varepsilon} \rangle_t = \int_0^t \langle d\Psi_f^\varepsilon, d\Psi_f^\varepsilon \rangle_{u/\varepsilon^2}(\mathbf{X}_u) du.$$

From (3.22), (3.20) and Burkholder-Davis-Gundy inequality, we see that there is a positive constant C such that

$$\begin{aligned} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left(\nabla \Psi_f^\varepsilon(\mathbf{X}_u), b\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) \right) du \right| \right] &\leq \varepsilon \mathbb{E}^\varepsilon [|\Psi_f^\varepsilon(\mathbf{X}_{t \wedge \tau}) - \Psi_f^\varepsilon(\mathbf{x})|] \\ &\quad + \varepsilon \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} (\mathcal{L}_{u/\varepsilon^2}^\varepsilon \Psi_f^\varepsilon)(\mathbf{X}_u) du \right| \right] + \varepsilon \mathbb{E}^\varepsilon \left[\left| \mathbf{M}_{t \wedge \tau}^{\Psi_f^\varepsilon, \varepsilon} \right| \right] \\ &< C \varepsilon^{1-p(n+1)} (1+t) \|f\|_{C^2(\bar{\mathbf{I}})}. \end{aligned} \quad (3.23)$$

Apply Lemma 3.1, with $(\nabla \Psi_f^\varepsilon, b)$ in place of f , and then recall (3.21) to notice that there exists a positive constant C such that

$$\begin{aligned} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((\nabla \Psi_f^\varepsilon(\mathbf{X}_u), b\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right)) - (\mathbb{M}(\nabla \Psi_f^\varepsilon, b))(\mathbf{X}_u) \right) du \right| \right] \\ &< C \varepsilon^{1-p(n+2)} (1+t) \|f\|_{C^3(\bar{\mathbf{I}})}. \end{aligned} \quad (3.24)$$

Finally, combine (3.23) and (3.24) into (3.15). $\square$

Lemma 3.3. *For any function $f \in C^3(\bar{\mathbf{I}} \times [0, \infty); \mathbb{R})$ which is 2π -periodic in its last argument and for any $0 < p < 1$ there exists a constant $C_c > 0$ such that*

$$\begin{aligned} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left(f\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{A}(\mathbb{M}f))([\mathbf{X}_u]) \right) \omega\left(\frac{K(\mathbf{X}_u)}{\varepsilon^p}\right) du \right| \right] \\ < C_c \varepsilon^{1-p(n+2)} (1+t) \|f\|_{C^3(\bar{\mathbf{I}} \times [0, \infty))}. \end{aligned}$$

Proof. We denote

$$q_\varepsilon(\mathbf{x}, t) := f(\mathbf{x}, t) \omega\left(\frac{K(\mathbf{x})}{\varepsilon^p}\right), \quad \mathbf{x} \in \bar{\mathbf{I}}, \quad t \geq 0.$$

Notice that $q_\varepsilon \in C^2(\bar{\mathbf{I}} \times [0, \infty); \mathbb{R})$ is 2π -periodic in its last argument,

$$\|q_\varepsilon\|_{C^2(\bar{\mathbf{I}} \times [0, \infty))} \leq C \varepsilon^{-2p} \|f\|_{C^2(\bar{\mathbf{I}} \times [0, \infty))} \quad (3.25)$$

for some constant $C > 0$, and

$$\begin{aligned} \left(f\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{A}(\mathbb{M}f))([\mathbf{X}_u]) \right) \omega\left(\frac{K(\mathbf{X}_u)}{\varepsilon^p}\right) &= \left(q_\varepsilon\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}q_\varepsilon)(\mathbf{X}_u) \right) \\ &+ \left((\mathbb{M}f)(\mathbf{X}_u) - (\mathbb{A}(\mathbb{M}f))([\mathbf{X}_u]) \right) \omega\left(\frac{K(\mathbf{X}_u)}{\varepsilon^p}\right). \end{aligned} \quad (3.26)$$

Applying Lemma 3.1, and using (3.25), we obtain

$$\mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left(q_\varepsilon\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}q_\varepsilon)(\mathbf{X}_u) \right) du \right| \right] < C \varepsilon^{1-2p} (1+t) \|f\|_{C^2(\bar{\mathbf{I}} \times [0, \infty))}. \quad (3.27)$$

The function $\mathbb{M}f \in C^3(\bar{\mathbf{I}}; \mathbb{R})$, and $\|\mathbb{M}f\|_{C^3(\bar{\mathbf{I}})} \leq \|f\|_{C^3(\bar{\mathbf{I}} \times [0, \infty))}$. Taking $\mathbb{M}f$ in place of f in Lemma 3.2 we have

$$\begin{aligned} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((\mathbb{M}f)(\mathbf{X}_u) - (\mathbb{A}(\mathbb{M}f))([\mathbf{X}_u]) \right) \omega\left(\frac{K(\mathbf{X}_u)}{\varepsilon^p}\right) du \right| \right] \\ < C_a \varepsilon^{1-p(n+2)} (1+t) \|f\|_{C^3(\bar{\mathbf{I}} \times [0, \infty))}. \end{aligned} \quad (3.28)$$

Combine (3.26), (3.27) and (3.28) to complete this proof. $\square$

Denote $\psi(\mathbf{x}, t) := (\nabla K(\mathbf{x}), b(\mathbf{x}, t))$, then $\mathbb{M}\psi \equiv 0$ and by definition (3.8),

$$\Phi_\psi(\mathbf{x}, t) = \int_0^t (\nabla K(\mathbf{x}), b(\mathbf{x}, u)) du.$$

Lemma 3.4. *For any function $g \in C^3(\mathbb{R}; \mathbb{R})$ and for any $0 < p < 1$ there exists a constant $C_b > 0$ such that*

$$\begin{aligned} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left\{ \frac{1}{\varepsilon} \psi\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) + (\nabla \Phi_\psi, b)\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) \right\} g(K(\mathbf{X}_u)) \omega\left(\frac{K(\mathbf{X}_u)}{\varepsilon^p}\right) du \right| \right] \\ < C_b \varepsilon^{1-3p} (1+t) \|g\|_{C^3([0, K^*])}. \end{aligned}$$

N. V. O'Bryant

Proof. Let $k_\varepsilon(h) = g(h) \omega(\frac{h}{\varepsilon^p})$, $h \in \mathbb{R}$. By the stopped martingale problem for $(\mathcal{L}_{t/\varepsilon^2}^\varepsilon, \delta_{\mathbf{x}}, \mathbf{I})$, $t \geq 0$,

$$\begin{aligned}
& \varepsilon \Phi_\psi \left(\mathbf{X}_{t \wedge \tau}, \frac{t \wedge \tau}{\varepsilon^2} \right) k_\varepsilon(K(\mathbf{X}_{t \wedge \tau})) - \varepsilon \Phi_\psi(\mathbf{x}, 0) k_\varepsilon(K(\mathbf{x})) \\
& - \frac{1}{\varepsilon} \int_0^{t \wedge \tau} \frac{\partial \Phi_\psi}{\partial u} \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) k_\varepsilon(K(\mathbf{X}_u)) du \\
& - \int_0^{t \wedge \tau} \left(\nabla \Phi_\psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right), b \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) \right) k_\varepsilon(K(\mathbf{X}_u)) du \\
& - \varepsilon \int_0^{t \wedge \tau} (\mathcal{L}_{u/\varepsilon^2} \Phi_\psi) \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) k_\varepsilon(K(\mathbf{X}_u)) du \\
& - \int_0^{t \wedge \tau} \Phi_\psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) \psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) k'_\varepsilon(K(\mathbf{X}_u)) du \\
& - \varepsilon \int_0^{t \wedge \tau} \Phi_\psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) (\mathcal{L}_{u/\varepsilon^2}(k_\varepsilon \circ K)) \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) du \\
& - 2\varepsilon \int_0^{t \wedge \tau} \langle d\Phi_\psi, d(k_\varepsilon \circ K) \rangle_{u/\varepsilon^2} \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) = \varepsilon M_{t \wedge \tau}^\varepsilon, \tag{3.29}
\end{aligned}$$

where M^ε is a $\mathbb{P}^\varepsilon$ -martingale with its quadratic variation

$$\langle M^\varepsilon \rangle_t = \int_0^t \langle d(\Phi_\psi(k_\varepsilon \circ K)), d(\Phi_\psi(k_\varepsilon \circ K)) \rangle_{u/\varepsilon^2} \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) du.$$

Now notice that

$$\frac{\partial \Phi_\psi}{\partial t}(\mathbf{x}, t) = \psi(\mathbf{x}, t),$$

rearrange the terms of (3.29) and apply Burkholder-Davis-Gundy inequality to see that there exists a constant $C > 0$, such that

$$\begin{aligned}
& \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left\{ \frac{1}{\varepsilon} \psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) + \left(\nabla \Phi_\psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right), b \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) \right) \right\} k_\varepsilon(K(\mathbf{X}_u)) du \right| \right] \\
& \leq \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \Phi_\psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) \psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) k'_\varepsilon(K(\mathbf{X}_u)) du \right| \right] \\
& \quad + C\varepsilon^{1-2p}(1+t)\|g\|_{C^2([0, K^*])}. \tag{3.30}
\end{aligned}$$

Next consider

$$(\mathbb{M}(\Phi_\psi \psi(k'_\varepsilon \circ K)))(\mathbf{x}) = k'_\varepsilon(K(\mathbf{x}))(\mathbb{M}(\Phi_\psi \psi))(\mathbf{x}).$$

Since $b(\mathbf{x}, 2\pi) = b(\mathbf{x}, 0)$ for any $\mathbf{x}$, we have $\Phi_\psi(\mathbf{x}, 2\pi) = \Phi_\psi(\mathbf{x}, 0)$ for any $\mathbf{x}$. Integration by parts reveals that

$$(\mathbb{M}(\Phi_\psi \psi))(\mathbf{x}) = \frac{1}{2\pi} \int_0^{2\pi} \Phi_\psi(\mathbf{x}, t) \frac{\partial \Phi_\psi}{\partial t}(\mathbf{x}, t) dt \equiv 0.$$

Writing $r_\varepsilon := \Phi_\psi \psi(k'_\varepsilon \circ K)$ we have $\mathbb{M}r_\varepsilon \equiv 0$. Apply Lemma 3.1 with r_ε in place of f to see that there exists $C > 0$ such that

$$\begin{aligned} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \Phi_\psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) \psi \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) k'_\varepsilon (K(\mathbf{X}_u)) du \right| \right] \\ = \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left(r_\varepsilon \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) - (\mathbb{M}r_\varepsilon)(\mathbf{X}_u) \right) du \right| \right] \\ < C\varepsilon^{1-3p} (1+t) \|g\|_{C^3([0, K^*])}. \end{aligned}$$

Combine this inequality with (3.30) to complete the proof. $\square$

3.3. Time spent near the boundary of the critical set

We will use Lemma 3.1 to show that, under $\mathbb{P}^\varepsilon$, the process $\mathbf{X}$ does not spend ‘too much time’ near the boundary $\partial \mathbf{V}$ of the critical set $\bar{\mathbf{V}}$. This fact will be effectively used later.

Lemma 3.5. *Given $t \geq 0$,*

$$\overline{\lim}_{\varepsilon \rightarrow 0} \overline{\lim}_{\delta \rightarrow 0} \mathbb{E}^\varepsilon \left[\frac{1}{\delta} \int_0^{t \wedge \tau} \chi_{[-\delta, \delta]}(K(\mathbf{X}_u)) du \right] < \infty.$$

Proof. Let $v \in C^\infty(\mathbb{R}; [0, 1])$ be an even function, such that

$$v(h) = \begin{cases} 1 & \text{if } |h| \leq 1, \\ 0 & \text{if } |h| \geq 2. \end{cases}$$

For every $h \in \mathbb{R}$ and $\delta > 0$, we define a function

$$g_\delta(h) := \frac{1}{\delta} \int_0^h \left(\int_0^r v\left(\frac{s}{\delta}\right) ds \right) dr,$$

so that

$$g''_\delta(h) = \frac{1}{\delta} v\left(\frac{h}{\delta}\right) \geq \frac{1}{\delta} \chi_{[-\delta, \delta]}(h) \quad (3.31)$$

and $g_\delta \in \mathcal{D}^\varepsilon$. Let $0 \leq t < \tau$ and $\mathbf{x} \in \bar{\mathbf{I}}$. By the martingale problem for $(\mathcal{L}_{t/\varepsilon^2}^\varepsilon, \delta_{\mathbf{x}}, \mathbf{I})$,

$$\begin{aligned} g_\delta(K(\mathbf{X}_t)) - g_\delta(K(\mathbf{x})) &- \frac{1}{\varepsilon} \int_0^t g'_\delta(K(\mathbf{X}_u)) \left(\nabla K(\mathbf{X}_u), b\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) \right) du \\ &- \int_0^t g'_\delta(K(\mathbf{X}_u)) (\mathcal{L}_{u/\varepsilon^2} K)(\mathbf{X}_u) du \\ &- \frac{1}{2} \int_0^t g''_\delta(K(\mathbf{X}_u)) \langle dK, dK \rangle_{u/\varepsilon^2}(\mathbf{X}_u) du \end{aligned}$$

N. V. O'Bryant

is a $\mathbb{P}^\varepsilon$ -martingale. Set

$$\begin{aligned} f_1(\mathbf{x}, t) &:= g'_\delta(K(\mathbf{x}))(\nabla K(\mathbf{x}), b(\mathbf{x}, t)), \\ f_2(\mathbf{x}, t) &:= g'_\delta(K(\mathbf{x}))(\mathcal{L}_t K)(\mathbf{x}), \\ f_3(\mathbf{x}, t) &:= g''_\delta(K(\mathbf{x}))\langle dK, dK \rangle_t(\mathbf{x}), \end{aligned}$$

so that

$$\begin{aligned} (\mathbb{M}f_1)(\mathbf{x}) &= g'_\delta(K(\mathbf{x}))(\nabla K(\mathbf{x}), (\mathbb{M}b)(\mathbf{x})) \\ &= nK^{n-1}(\mathbf{x})g'_\delta(K(\mathbf{x}))(\nabla K(\mathbf{x}), \nabla^\perp K(\mathbf{x})) \equiv 0, \\ (\mathbb{M}f_2)(\mathbf{x}) &= g'_\delta(K(\mathbf{x}))(\mathbb{M}(\mathcal{L}K))(\mathbf{x}), \\ (\mathbb{M}f_3)(\mathbf{x}) &= g''_\delta(K(\mathbf{x}))(\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}), \end{aligned}$$

and

$$\begin{aligned} &g_\delta(K(\mathbf{X}_t)) - g_\delta(K(\mathbf{x})) \\ &\quad - \frac{1}{\varepsilon} \int_0^t \left(f_1\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}f_1)(\mathbf{X}_u) \right) du \\ &\quad - \int_0^t \left(f_2\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}f_2)(\mathbf{X}_u) \right) du \\ &\quad - \frac{1}{2} \int_0^t \left(f_3\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}f_3)(\mathbf{X}_u) \right) du \\ &\quad - \int_0^t (\mathbb{M}f_2)(\mathbf{X}_u) du \\ &\quad - \frac{1}{2} \int_0^t (\mathbb{M}f_3)(\mathbf{X}_u) du \end{aligned}$$

is a $\mathbb{P}^\varepsilon$ -martingale. Then for any $\varepsilon > 0, \delta > 0$, and for any $0 \leq t \leq \tau$,

$$\begin{aligned} \frac{1}{2} \mathbb{E}^\varepsilon \left[\int_0^t (\mathbb{M}f_3)(\mathbf{X}_u) du \right] &\leq \mathbb{E}^\varepsilon \left[\left| g_\delta(K(\mathbf{X}_t)) - g_\delta(K(\mathbf{x})) \right| \right] \\ &\quad + \frac{1}{\varepsilon} \mathbb{E}^\varepsilon \left[\left| \int_0^t \left(f_1\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}f_1)(\mathbf{X}_u) \right) du \right| \right] \\ &\quad + \mathbb{E}^\varepsilon \left[\left| \int_0^t \left(f_2\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}f_2)(\mathbf{X}_u) \right) du \right| \right] \\ &\quad + \frac{1}{2} \mathbb{E}^\varepsilon \left[\left| \int_0^t \left(f_3\left(\mathbf{X}_u, \frac{u}{\varepsilon^2}\right) - (\mathbb{M}f_3)(\mathbf{X}_u) \right) du \right| \right] \\ &\quad + \mathbb{E}^\varepsilon \left[\left| \int_0^t (\mathbb{M}f_2)(\mathbf{X}_u) du \right| \right]. \end{aligned} \tag{3.32}$$

We use Remark 2.1 to make sure that

$$C_1 := \frac{1}{2} \inf_{\mathbf{y} \in \bar{\mathcal{E}}} (\mathbb{M}\langle dK, dK \rangle)(\mathbf{y}) > 0,$$

which, together with inequality (3.31), guarantees that for all $\mathbf{x} \in \bar{\mathbf{I}}$ and for any $0 < \delta < a$

$$(\mathbb{M}f_3)(\mathbf{x}) \geq C_1 \frac{1}{\delta} \chi_{[-\delta, \delta]}(K(\mathbf{x})).$$

Notice that each of the functions $f_i \in C^2(\bar{\mathbf{I}} \times [0, \infty); \mathbb{R})$, $i = 1, 2, 3$, is 2π -periodic in its last argument, i.e., each of these three functions f_i satisfy all the hypotheses on the function f in Lemma 3.1. Now, inequality (3.32) and Lemma 3.1 together imply that for all $\mathbf{x} \in \bar{\mathbf{I}}$, all $0 \leq t \leq \tau$, and for all $0 < \delta < a$,

$$\begin{aligned} C_1 \mathbb{E}^\varepsilon \left[\frac{1}{\delta} \int_0^t \chi_{[-\delta, \delta]}(K(\mathbf{X}_u)) du \right] &\leq \mathbb{E}^\varepsilon \left[\left| g_\delta(K(\mathbf{X}_t)) - g_\delta(K(\mathbf{x})) \right| \right] \\ &+ C(1+t) \left(\|f_1\|_{C^2(\bar{\mathbf{I}} \times [0, \infty))} + \varepsilon \|f_2\|_{C^2(\bar{\mathbf{I}} \times [0, \infty))} + \varepsilon \|f_3\|_{C^2(\bar{\mathbf{I}} \times [0, \infty))} \right) \\ &+ \mathbb{E}^\varepsilon \left[\left| \int_0^t (\mathbb{M}f_2)(\mathbf{X}_u) du \right| \right]. \quad (3.33) \end{aligned}$$

For all $\delta > 0$, there exists a constant $C^* > 0$ such that

$$\sup_{-K^* \leq h \leq K^*} |g'_\delta(h)| \leq C^* \quad \text{and} \quad \sup_{-K^* \leq h \leq K^*} |g_\delta(h)| \leq C^*.$$

Then, for all $\delta > 0$, all $0 \leq t \leq \tau$, and $\mathbf{x} \in \bar{\mathbf{I}}$,

$$\mathbb{E}^\varepsilon \left[\left| g_\delta(K(\mathbf{X}_t)) - g_\delta(K(\mathbf{x})) \right| \right] \leq 2C^* \quad (3.34)$$

and there exists a positive constant C^{**} such that

$$\mathbb{E}^\varepsilon \left[\left| \int_0^t (\mathbb{M}f_2)(\mathbf{X}_u) du \right| \right] < C^{**} t \|K\|_{C^2(\bar{\mathbf{I}})}. \quad (3.35)$$

Denote $C_{f_i} = \|f_i\|_{C^2(\bar{\mathbf{I}} \times [0, \infty))}$, $i = 1, 2, 3$, $C_K = \|K\|_{C^2(\bar{\mathbf{I}})}$. Plug inequalities (3.34) and (3.35) into (3.33) to obtain that for all $t > 0$,

$$\begin{aligned} \overline{\lim}_{\delta \rightarrow 0} \mathbb{E}^\varepsilon \left[\frac{1}{\delta} \int_0^{t \wedge \tau} \chi_{[-\delta, \delta]}(K(\mathbf{X}_u)) du \right] \\ \leq \frac{1}{C_1} (2C^* + C(1+t)(C_{f_1} + \varepsilon C_{f_2} + \varepsilon C_{f_3}) + C^{**} C_K t). \end{aligned}$$

Finally, let ε tend to 0 to see that the last inequality implies that

$$\overline{\lim}_{\varepsilon \rightarrow 0} \overline{\lim}_{\delta \rightarrow 0} \mathbb{E}^\varepsilon \left[\frac{1}{\delta} \int_0^{t \wedge \tau} \chi_{[-\delta, \delta]}(K(\mathbf{X}_u)) du \right] \leq \overline{C} + \overline{\overline{C}} t,$$

where $\overline{C} = (2C^* + C C_{f_1}) / C_1 < \infty$, and $\overline{\overline{C}} = (C^{**} C_K + C C_{f_1}) / C_1 < \infty$. $\square$

Remark 3.4. For the purpose of our proof, the lemma above is not stated in the most general form. The following more general result holds.

N. V. O'Bryant

Given $t \geq 0$, and a function $v \in L_1(\mathbb{R})$,

$$\overline{\lim}_{\varepsilon \rightarrow 0} \overline{\lim}_{\delta \rightarrow 0} \mathbb{E}^\varepsilon \left[\frac{1}{\delta} \int_0^{t \wedge \tau} v \left(\frac{K(\mathbf{X}_u)}{\delta} \right) du \right] < \infty. \quad (3.36)$$

We can prove this statement analogously, setting $g'_\delta(h) = \frac{1}{\delta} v\left(\frac{h}{\delta}\right)$. Then $g'_\delta(h) = \int_0^{h/\delta} v(s) ds$, and $g_\delta(h) = \delta \int_0^{h/\delta} \left(\int_0^s v(u) du \right) ds$.

For all $\delta > 0$, there exists a constant $C^* > 0$ such that

$$\sup_{-K^* \leq h \leq K^*} |g'_\delta(h)| \leq \int_0^\infty v(s) ds \leq C^*$$

and

$$\sup_{-K^* \leq h \leq K^*} |g_\delta(h)| \leq \sup_{-K^* \leq h \leq K^*} \left| \delta \int_0^{h/\delta} \left(\int_0^\infty v(u) du \right) ds \right| \leq C^*.$$

Now, to verify inequality (3.36), we can repeat all the arguments of the proof of Lemma 3.5.

3.4. Stochastic averaging around the boundary of the critical set

In order to perform analysis close to $\partial \mathbf{V}$, and to achieve sufficient accuracy, we will introduce new coordinates defined on $\mathcal{E}$. We will use the flow ζ_t defined on $\mathcal{E}$ by

$$\dot{\zeta}_t(\mathbf{x}) = \frac{\nabla K}{\|\nabla K\|^2}(\zeta_t(\mathbf{x})), \quad \zeta_0(\mathbf{x}) = \mathbf{x}. \quad (3.37)$$

Notice that for any $\mathbf{x} \in \mathcal{E}$

$$\frac{dK}{dt}(\zeta_t(\mathbf{x})) = \left(\nabla K(\zeta_t(\mathbf{x})), \dot{\zeta}_t(\mathbf{x}) \right) \equiv 1,$$

and if $\mathbf{x} \in \partial \mathbf{V}$, then $K(\zeta_t(\mathbf{x})) = t$ for all $-a < t < a$. Fix a point $\mathbf{x}^* \in \partial \mathbf{V}$, and define the curve

$$\mathcal{C} := \{\zeta_{-s}(\mathbf{x}^*) : -a < s < a\}.$$

We reverse time to emphasize that a trajectory of the constructed flow moves a point $\mathbf{x} \in \mathcal{E} \cup \mathcal{C}$ towards (not away from) $\mathbf{x}^* \in \partial \mathbf{V}$.

Notice that $\mathcal{E} = \{\phi_t(\mathbf{x}) : \mathbf{x} \in \mathcal{C}, 0 \leq t < \eta(K(\mathbf{x}))\}$, and in particular the map $(t, s) \mapsto \phi_t(\zeta_s(\mathbf{x}^*))$ is a homeomorphism from

$$\{(t, s) \in \mathbb{R}^2 : -a < s < a, 0 \leq t < \eta(K(\zeta_s(\mathbf{x}^*)))\}$$

to $\mathcal{E}$. This is the foundation of our new coordinate system for $\mathcal{E}$. It is advantageous to normalize the first coordinate by setting

$$\beta(s) := \int_{\mathbf{y} \in K^{-1}(s)} \frac{(\mathbb{M}\langle dK, dK \rangle)(\mathbf{y})}{\|\nabla K(\mathbf{y})\|} dl(y), \quad \Theta(\mathbf{x}) := \int_0^t (\mathbb{M}\langle dK, dK \rangle)(\phi_s(\mathbf{x})) ds,$$

where t satisfies $\phi_t(\zeta_s(\mathbf{x}^*)) = \mathbf{x}$. The map $\rho : \mathbf{x} \mapsto (\Theta(\mathbf{x}), K(\mathbf{x}))$ is a homeomorphism from the set $\mathcal{E}$ onto $\{(\theta, s) \in \mathbb{R}^2 : -a < s < a, 0 \leq \theta < \beta(s)\}$. For future reference, notice that for any $\mathbf{x} \in \mathcal{E} \setminus \mathcal{C}$

$$(\nabla \Theta(\mathbf{x}), \nabla^\perp K(\mathbf{x})) = (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}). \quad (3.38)$$

This equation reflects the essential property of our new coordinates, that equal increments of Θ correspond to equal amounts of diffusion across $\partial \mathbf{V}$. Recall the definition (3.14) of Ψ_f , and define

$$\check{\Psi}_f := \Psi_f \circ \rho^{-1}.$$

That is, for any $\mathbf{x} \in \mathcal{E}$, $\Psi_f(\mathbf{x}) = \check{\Psi}_f(\theta, s)$ with $\theta = \Theta(\mathbf{x})$ and $s = K(\mathbf{x})$. We can specify f so that $\check{\Psi}_f(\theta, 0) \in C^5(\mathbb{R})$, and $\check{\Psi}_f(\theta + \beta(s), s) = \check{\Psi}_f(\theta, s)$, $\theta \in \mathbb{R}$, $-a < s < a$. Using property (3.38) of the new coordinates, for all $\mathbf{x} \in \mathcal{E} \setminus \mathcal{C}$,

$$\begin{aligned} (\nabla \Psi_f(\mathbf{x}), \nabla^\perp K(\mathbf{x})) &= \left(\frac{\partial \check{\Psi}_f}{\partial \theta} \nabla \Theta(\mathbf{x}) + \frac{\partial \check{\Psi}_f}{\partial s} \nabla K(\mathbf{x}), \nabla^\perp K(\mathbf{x}) \right) \\ &= \frac{\partial \check{\Psi}_f}{\partial \theta} \left(\nabla \Theta(\mathbf{x}), \nabla^\perp K(\mathbf{x}) \right) = \frac{\partial \check{\Psi}_f}{\partial \theta} (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}), \end{aligned} \quad (3.39)$$

with all the derivatives of $\check{\Psi}_f$ computed at $(\Theta(\mathbf{x}), K(\mathbf{x}))$.

As before, we assume that function $\omega \in C_c^\infty(\mathbb{R})$ is supported on a compact subset of $(0, \infty)$. We will borrow the following proposition from [?], only with minor changes.

Proposition 3.2. *Let $p = 1/(n+1)$ and*

$$\overline{\omega}(h) := \omega(h^{2p}) h^{2p}, \quad h \in \mathbb{R}.$$

There exists a function $B \in C^2(\mathbb{R} \times [0, \infty))$ such that

$$\begin{aligned} n \frac{\partial B}{\partial \theta} + \frac{(n+1)^2}{8} \frac{\partial^2 B}{\partial s^2} + \frac{n^2-1}{8s} \frac{\partial B}{\partial s} &= \frac{\partial \check{\Psi}_f(\theta, 0)}{\partial \theta} \overline{\omega}(s), \\ B(\theta + \beta(0), s) &= B(\theta, s), \\ |\partial^\alpha B(\theta, s)| &\leq C \left\| \check{\Psi}_f(\cdot, 0) \right\|_{C^5(\mathbb{R})} e^{-s/C} \end{aligned} \quad (3.40)$$

for all $(\theta, s) \in \mathbb{R} \times (0, \infty)$, for some constant $C > 0$, and for any multi-index α , $|\alpha| \leq 2$, and such that

$$\frac{\partial B(\theta, 0)}{\partial s} = 0 \quad (3.41)$$

for all $\theta \in \mathbb{R}$.

It was shown in [?] that solving (3.40) involves Fourier analysis and leads to an inhomogeneous Bessel's ordinary differential equation. The unique explicit solution is expressed via Bessel functions of small order and purely imaginary argument.

Define the functions $\mathbf{d}_n : \mathbb{R} \rightarrow \mathbb{R}$ by $\mathbf{d}_n(h) := \left(\max\{0, h\} \right)_n$. This sequence satisfies the recursive relation $\mathbf{d}_n(h) = h \mathbf{d}_{n-1}(h)$. For some $n > 2$, $H(\mathbf{x}) = \mathbf{d}_n(K(\mathbf{x}))$ and $\nabla^\perp H(\mathbf{x}) = n \mathbf{d}_{n-1}(K(\mathbf{x})) \nabla^\perp K(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^2$.

N. V. O'Bryant

Proposition 3.3. *There exists a constant $C > 0$ such that for any $f \in C^5(\mathbb{R}^2; \mathbb{R})$ and for any $0 < p < 1$*

$$\left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} \left(f(\mathbf{X}_u) - (\mathbb{A}f)([\mathbf{X}_u]) \right) d_n(K(\mathbf{X}_u)) \omega \left(\frac{K(\mathbf{X}_u)}{\varepsilon^p} \right) du \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right| < C\varepsilon^{1+p} (1+t) \|f\|_{C^5(\bar{\mathbf{I}})}. \quad (3.42)$$

whenever $0 \leq t_1 < t_2 < \dots < t_m \leq s < t$, and $h_1, \dots, h_m \in C_b(\mathbb{R}^2)$.

Remark 3.5. Since ω has compact support we can find a constant $C^* > 0$ such that for any $\mathbf{x} \in \bar{\mathbf{I}}$

$$\left| d_n(K(\mathbf{x})) \omega \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \right| < C^* \varepsilon^{np} \chi_{[-\varepsilon^p, \varepsilon^p]}(K(\mathbf{x})).$$

Then, by Lemma 3.5, there exist constants $C^{**}, C > 0$, such that

$$\begin{aligned} \left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} \left(f(\mathbf{X}_u) - (\mathbb{A}f)([\mathbf{X}_u]) \right) d_n(K(\mathbf{X}_u)) \omega \left(\frac{K(\mathbf{X}_u)}{\varepsilon^p} \right) du \right] \right| \\ < C^{**} \varepsilon^{np} \|f\|_{C(\bar{\mathbf{I}})} \mathbb{E}^\varepsilon \int_{s \wedge \tau}^{t \wedge \tau} \chi_{[-\varepsilon^p, \varepsilon^p]}(K(\mathbf{X}_u)) du \\ < C \varepsilon^{p(n+1)} \|f\|_{C(\bar{\mathbf{I}})}. \end{aligned}$$

For values of $p < 1/n$, Proposition 3.3 provides a better bound.

Proof. For every $\mathbf{x} \in \bar{\mathbf{I}}$ and for all $f \in C^4(\mathbb{R}^2; \mathbb{R})$ we define

$$\mathbb{G}_f^\varepsilon(\mathbf{x}) := \frac{1}{\varepsilon} \left(f(\mathbf{x}) - (\mathbb{A}_0 f)([\mathbf{x}]) \right) d_n(K(\mathbf{x})) \omega \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right).$$

We will perform stochastic averaging using the martingale problem. This approach was developed by Papanicolaou and Kohler [?]. We will look for a function $\Psi^\varepsilon \in C^2(\mathbb{R}^2)$ that satisfies

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{-p} \|\Psi^\varepsilon\|_{C(\bar{\mathbf{I}})} < \infty,$$

and which, for every $\varepsilon > 0, t \geq 0$, solves the approximate partial differential equation

$$(\mathbb{M}(\mathcal{L}^\varepsilon \Psi^\varepsilon))(\mathbf{x}) \approx \mathbb{G}_f^\varepsilon(\mathbf{x}), \quad \mathbf{x} \in \bar{\mathbf{I}}. \quad (3.43)$$

From the approximate equation (3.43), by Lemma 3.1, and by the stopped martingale problem for $(\mathcal{L}_{t/\varepsilon^2}^\varepsilon, \delta_{\mathbf{x}}, \mathbf{I})$,

$$\begin{aligned} \int_0^{t \wedge \tau} \mathbb{G}_f^\varepsilon(\mathbf{X}_u) du &\approx \int_0^{t \wedge \tau} (\mathbb{M}(\mathcal{L}^\varepsilon \Psi^\varepsilon))(\mathbf{X}_u) du \approx \int_0^{t \wedge \tau} (\mathcal{L}_{u/\varepsilon^2}^\varepsilon \Psi^\varepsilon)(\mathbf{X}_u) du \\ &= \Psi^\varepsilon(\mathbf{X}_{t \wedge \tau}) - \Psi^\varepsilon(\mathbf{X}_0) + M_{t \wedge \tau}^\varepsilon, \end{aligned}$$

where $M_{t \wedge \tau}^\varepsilon$ is a $\mathbb{P}^\varepsilon$ -martingale. Then, we will finish the proof by optional sampling for this martingale.

The operator

$$\varepsilon \mathbb{M}(\mathcal{L}^\varepsilon \Psi^\varepsilon) = (\nabla \Psi^\varepsilon(\mathbf{x}), \nabla^\perp H(\mathbf{x})) + \varepsilon \mathbb{M}(\mathcal{L} \Psi^\varepsilon)$$

has drift of order 1 and small diffusion, which leads to a singular perturbation problem.

Notice from (3.19) and from property (3.39) of the new coordinates that for every $\mathbf{x} \in \mathcal{E} \setminus \mathcal{C}$

$$\mathbb{G}_f^\varepsilon(\mathbf{x}) := \varepsilon^{-1} \frac{\partial \check{\Psi}_f(\Theta(\mathbf{x}), K(\mathbf{x}))}{\partial \theta} \mathbf{d}_n(K(\mathbf{x})) \omega\left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}). \quad (3.44)$$

Recall that $\overline{\omega}(h) = \omega(h^{2p}) h^{2p}$, $h \in \mathbb{R}$, and write $\mathbf{d}_n(h)$ as $\mathbf{d}_{n-1}(h)h$. Then, from (3.44),

$$\mathbb{G}_f^\varepsilon(\mathbf{x}) := \varepsilon^{p-1} \frac{\partial \check{\Psi}_f(\Theta(\mathbf{x}), K(\mathbf{x}))}{\partial \theta} \mathbf{d}_{n-1}(K(\mathbf{x})) \overline{\omega}\left(\frac{\mathbf{d}_{1/2p}(K(\mathbf{x}))}{\sqrt{\varepsilon}}\right) (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}).$$

For all $\mathbf{x} \in \mathcal{E}$ we define

$$\Theta^\circ(\mathbf{x}) := \frac{\beta(0)}{\beta(K(\mathbf{x}))} \Theta(\mathbf{x}).$$

To continue with this proof, we state and prove the following lemma.

Lemma 3.6. *Let $f \in C^5(\mathbb{R}^2; \mathbb{R})$. Let B satisfy the hypotheses of Proposition 3.2, and for every $\varepsilon > 0$ define*

$$B^\varepsilon(\mathbf{x}) := \varepsilon^p B\left(\Theta^\circ(\mathbf{x}), \frac{\mathbf{d}_{1/2p}(K(\mathbf{x}))}{\sqrt{\varepsilon}}\right)$$

for all $\mathbf{x} \in \mathcal{E}$. Then

$$B^\varepsilon \in C^1(\mathcal{E}) \cap C^2(\mathcal{E} \setminus (\partial \mathbf{V} \cup \mathcal{C})),$$

and

$$(\mathbb{M}(\mathcal{L}^\varepsilon B^\varepsilon))(\mathbf{x}) := \lim_{\substack{\mathbf{y} \rightarrow \mathbf{x} \\ \mathbf{y} \in \mathcal{E} \setminus (\partial \mathbf{V} \cup \mathcal{C})}} (\mathbb{M}(\mathcal{L}^\varepsilon B^\varepsilon))(\mathbf{y})$$

exists for all $\mathbf{x} \in \partial \mathbf{V} \cup \mathcal{C}$. There exist positive constants $C_1, C_2 > 0$ such that

$$|(\mathbb{M}(\mathcal{L}^\varepsilon B^\varepsilon))(\mathbf{x}) - \mathbb{G}_f^\varepsilon(\mathbf{x})| \leq C_1 \|f\|_{C^5(\bar{\mathbf{I}})} \exp\left\{-\frac{|K(\mathbf{x})|^{1/2p}}{C_2 \sqrt{\varepsilon}}\right\},$$

$$\sqrt{(\mathbb{M}\langle dB^\varepsilon, dB^\varepsilon \rangle)(\mathbf{x})} \leq C_1 \|f\|_{C^5(\bar{\mathbf{I}})} \exp\left\{-\frac{|K(\mathbf{x})|^{1/2p}}{C_2 \sqrt{\varepsilon}}\right\},$$

and

$$|B^\varepsilon(\mathbf{x})| \leq C_1 \varepsilon^p \|f\|_{C^5(\bar{\mathbf{I}})}$$

for all $\mathbf{x} \in \mathcal{E} \setminus (\partial \mathbf{V} \cup \mathcal{C})$ and $\varepsilon > 0$.

N. V. O'Bryant

Proof. First of all, recall that the small set $\mathcal{E} = \{\mathbf{y} \in \mathbb{R}^2 : |K(\mathbf{y})| < a\} \supset \partial\mathbf{V}$ is homeomorphic to the set $\{(\theta, s) \in \mathbb{R}^2 : -a < s < a, 0 \leq \theta < \beta(s)\}$, and that $\mathcal{C}$ is a fixed curve in $\mathcal{E}$, which crosses all $K^{-1}(s)$ transversally.

Notice that the definition of B^ε can be rewritten more explicitly as

$$B^\varepsilon(\mathbf{x}) := \begin{cases} \varepsilon^p B(\Theta^\circ(\mathbf{x}), 0) & \text{if } -a < K(\mathbf{x}) < 0, \\ \varepsilon^p B\left(\Theta^\circ(\mathbf{x}), \sqrt{\frac{K^{n+1}(\mathbf{x})}{\varepsilon}}\right) & \text{if } 0 \leq K(\mathbf{x}) < a. \end{cases}$$

We know that $B \in C^2(\mathbb{R} \times [0, \infty))$ is periodic in its first argument, which, together with the properties of Θ° and $d_{1/2p} \circ K$ away from $\partial\mathbf{V} \cup \mathcal{C}$, guarantees that $B^\varepsilon \in C^2(\mathcal{E} \setminus (\partial\mathbf{V} \cup \mathcal{C}))$.

The function K is smooth inside $\mathcal{E}$, and $1/2p = (n-1)/2 > 0$ implies differentiability of $d_{1/2p}$ on $(-a, a)$. Therefore, we are guaranteed that $d_{1/2p} \circ K \in C^1(\mathcal{E})$. By the definition of Θ° , it is guaranteed to be differentiable on $\mathcal{E}$. Notice that given any $\mathbf{x} \in \mathcal{E}$, we have $\text{span}\{\nabla K(\mathbf{x}), \nabla^\perp K(\mathbf{x})\} = \mathbb{R}^2$. From the periodicity in the first argument and from the boundary conditions on B , we also see that for every $\mathbf{x} \in \mathcal{E}$ both $\lim_{\mathbf{y} \rightarrow \mathbf{x}} (\nabla K, \nabla B^\varepsilon)(\mathbf{y})$ and $\lim_{\mathbf{y} \rightarrow \mathbf{x}} (\nabla^\perp K, \nabla B^\varepsilon)(\mathbf{y})$ exist. Therefore, $B^\varepsilon \in C^1(\mathcal{E})$. Thus, $B^\varepsilon \in C^1(\mathcal{E}) \cap C^2(\mathcal{E} \setminus (\partial\mathbf{V} \cup \mathcal{C}))$.

Let $\mathbf{x} \in \mathcal{E} \setminus (\partial\mathbf{V} \cup \mathcal{C})$, and consider

$$\begin{aligned} \mathcal{L}_t^\varepsilon B\left(\Theta^\circ(\mathbf{x}), \frac{d_{1/2p}(K(\mathbf{x}))}{\sqrt{\varepsilon}}\right) &= \frac{1}{2} \frac{\partial^2 B}{\partial \theta^2} \langle d\Theta^\circ, d\Theta^\circ \rangle_t(\mathbf{x}) + \frac{\partial B}{\partial \theta} (\mathcal{L}_t \Theta^\circ)(\mathbf{x}) \\ &\quad + \frac{1}{\sqrt{\varepsilon}} d'_{1/2p}(K(\mathbf{x})) \frac{\partial^2 B}{\partial s \partial \theta} \langle dK, d\Theta^\circ \rangle_t(\mathbf{x}) \\ &\quad + \frac{1}{2\varepsilon} \frac{\partial^2 B}{\partial s^2} \left(d'_{1/2p}(K(\mathbf{x}))\right)^2 \langle dK, dK \rangle_t(\mathbf{x}) \\ &\quad + \frac{1}{2\sqrt{\varepsilon}} \frac{\partial B}{\partial s} d''_{1/2p}(K(\mathbf{x})) \langle dK, dK \rangle_t(\mathbf{x}) \\ &\quad + \frac{1}{\sqrt{\varepsilon}} \frac{\partial B}{\partial s} d'_{1/2p}(K(\mathbf{x})) (\mathcal{L}_t K)(\mathbf{x}), \end{aligned} \quad (3.45)$$

and

$$\begin{aligned} \langle dB, dB \rangle_t(\mathbf{x}) &= \left(\frac{\partial B}{\partial \theta}\right)^2 \langle d\Theta^\circ, d\Theta^\circ \rangle_t(\mathbf{x}) \\ &\quad + \frac{1}{\varepsilon} \left(\frac{\partial B}{\partial s}\right)^2 \left(d'_{1/2p}(K(\mathbf{x}))\right)^2 \langle dK, dK \rangle_t(\mathbf{x}) \\ &\quad + \frac{2}{\sqrt{\varepsilon}} \frac{\partial B}{\partial \theta} \frac{\partial B}{\partial s} d'_{1/2p}(K(\mathbf{x})) \langle d\Theta^\circ, dK \rangle_t(\mathbf{x}). \end{aligned} \quad (3.46)$$

Here and below, B and its partial derivatives are evaluated at $\left(\Theta^\circ(\mathbf{x}), \frac{d_{1/2p}(K(\mathbf{x}))}{\sqrt{\varepsilon}}\right)$, unless specified otherwise. From (3.45) and (3.46), using

$$d'_{1/2p}(h) = \frac{1}{2p} d_{1/2p-1}(h)$$

and

$$\mathbf{d}_{1/2p}''(h) = \frac{1}{2p} \left(\frac{1}{2p} - 1 \right) \mathbf{d}_{1/2p-2}(h),$$

we can see that

$$\begin{aligned} (\mathcal{L}_t^\varepsilon B^\varepsilon)(\mathbf{x}) &= \frac{1}{\varepsilon} (\nabla B^\varepsilon(\mathbf{x}), b(\mathbf{x}, t)) + (\mathcal{L}_t B^\varepsilon)(\mathbf{x}) = \varepsilon^{p-1} \frac{\partial B}{\partial \theta} (\nabla \Theta^\circ(\mathbf{x}), b(\mathbf{x}, t)) \\ &\quad + \varepsilon^{p-3/2} \frac{\partial B}{\partial s} m \mathbf{d}_{1/2p-1}(K(\mathbf{x})) (\nabla K(\mathbf{x}), b(\mathbf{x}, t)) \\ &\quad + \varepsilon^p \left(\frac{1}{2} \frac{\partial^2 B}{\partial \theta^2} \langle d\Theta^\circ d\Theta^\circ \rangle_t(\mathbf{x}) + \frac{\partial B}{\partial \theta} (\mathcal{L}_t \Theta^\circ)(\mathbf{x}) \right. \\ &\quad + \frac{1}{\sqrt{\varepsilon}} \frac{1}{2p} \mathbf{d}_{1/2p-1}(K(\mathbf{x})) \frac{\partial^2 B}{\partial s \partial \theta} \langle dK, d\Theta^\circ \rangle_t(\mathbf{x}) \\ &\quad + \frac{1}{2\varepsilon} \frac{\partial^2 B}{\partial s^2} \left(\frac{1}{2p} \right)^2 \mathbf{d}_{1/2p-1}^2(K(\mathbf{x})) \langle dK, dK \rangle_t(\mathbf{x}) \\ &\quad + \frac{1}{2\sqrt{\varepsilon}} \frac{\partial B}{\partial s} \frac{1}{2p} \left(\frac{1}{2p} - 1 \right) \mathbf{d}_{1/2p-2}(K(\mathbf{x})) \langle dK, dK \rangle_t(\mathbf{x}) \\ &\quad \left. + \frac{1}{\sqrt{\varepsilon}} \frac{\partial B}{\partial s} \frac{1}{2p} \mathbf{d}_{1/2p-1}(K(\mathbf{x})) (\mathcal{L}_t K)(\mathbf{x}) \right) \end{aligned} \quad (3.47)$$

and

$$\begin{aligned} \langle dB^\varepsilon, dB^\varepsilon \rangle_t(\mathbf{x}) &= \varepsilon^{2p} \left(\frac{\partial B}{\partial \theta} \right)^2 \langle d\Theta^\circ, d\Theta^\circ \rangle_t(\mathbf{x}) \\ &\quad + \varepsilon^{2p-1} \left(\frac{1}{2p} \right)^2 \left(\frac{\partial B}{\partial s} \right)^2 \mathbf{d}_{1/2p-1}^2(K(\mathbf{x})) \langle dK, dK \rangle_t(\mathbf{x}) \\ &\quad + 2\varepsilon^{2p-1/2} \frac{\partial B}{\partial \theta} \frac{\partial B}{\partial s} \frac{1}{2p} \mathbf{d}_{1/2p-1}(K(\mathbf{x})) \langle dK, d\Theta^\circ \rangle_t(\mathbf{x}). \end{aligned} \quad (3.48)$$

Recall that $(\mathbb{M}(\nabla K, b))(\mathbf{x}) = 0$. Then, from (3.47),

$$\begin{aligned} (\mathbb{M}(\mathcal{L}^\varepsilon B^\varepsilon))(\mathbf{x}) &= \frac{1-2p}{8p^2} \varepsilon^{p-1/2} \mathbf{d}_{1/2p-2}(K(\mathbf{x})) \frac{\partial B}{\partial s} (\mathbb{M} \langle dK, dK \rangle)(\mathbf{x}) \\ &\quad + \frac{1}{8p^2} \varepsilon^{p-1} \mathbf{d}_{1/2p-1}^2(K(\mathbf{x})) \frac{\partial^2 B}{\partial s^2} (\mathbb{M} \langle dK, dK \rangle)(\mathbf{x}) \\ &\quad + \frac{1}{2p} \varepsilon^{p-1/2} \mathbf{d}_{1/2p-1}(K(\mathbf{x})) \frac{\partial B}{\partial s} (\mathbb{M}(\mathcal{L}K))(\mathbf{x}) \\ &\quad + \frac{1}{2p} \varepsilon^{p-1/2} \mathbf{d}_{1/2p-1}(K(\mathbf{x})) \frac{\partial^2 B}{\partial s \partial \theta} (\mathbb{M} \langle dK, d\Theta^\circ \rangle)(\mathbf{x}) \\ &\quad + \varepsilon^p \left[\frac{\partial B}{\partial \theta} (\mathbb{M}(\mathcal{L}\Theta^\circ))(\mathbf{x}) + \frac{1}{2} \frac{\partial^2 B}{\partial \theta^2} (\mathbb{M} \langle d\Theta^\circ, d\Theta^\circ \rangle)(\mathbf{x}) \right] \\ &\quad + n\varepsilon^{p-1} \mathbf{d}_{n-1}(K(\mathbf{x})) \frac{\partial B}{\partial \theta} (\nabla \Theta^\circ(\mathbf{x}), \nabla^\perp K(\mathbf{x})), \end{aligned}$$

N. V. O'Bryant

and from (3.48),

$$\begin{aligned} (\mathbb{M} \langle dB^\varepsilon, dB^\varepsilon \rangle) (\mathbf{x}) &= \varepsilon^{2p} \left(\frac{\partial B}{\partial \theta} \right)^2 (\mathbb{M} \langle d\Theta^\circ, d\Theta^\circ \rangle) (\mathbf{x}) \\ &\quad + \frac{1}{4p^2} \varepsilon^{2p-1} \left(\frac{\partial B}{\partial s} \right)^2 d_{1/2p-1}^2 (K(\mathbf{x})) (\mathbb{M} \langle dK, dK \rangle) (\mathbf{x}) \\ &\quad + \frac{1}{p} \varepsilon^{2p-1/2} \frac{\partial B}{\partial \theta} \frac{\partial B}{\partial s} d_{1/2p-1} (K(\mathbf{x})) (\mathbb{M} \langle dK, d\Theta^\circ \rangle) (\mathbf{x}). \end{aligned}$$

From (3.38),

$$(\nabla \Theta^\circ(\mathbf{x}), \nabla^\perp K(\mathbf{x})) = \frac{\beta(0)}{\beta(K(\mathbf{x}))} (\mathbb{M} \langle dK, dK \rangle) (\mathbf{x}).$$

Take $p = \frac{1}{n+1}$. Then, $\frac{1-2p}{8p^2} = \frac{n^2-1}{8}$, $\frac{1}{8p^2} = \frac{(n+1)^2}{8}$, and

$$\begin{aligned} (\mathbb{M} (\mathcal{L}^\varepsilon B^\varepsilon)) (\mathbf{x}) &= \varepsilon^{p-1} d_{n-1} (K(\mathbf{x})) (\mathbb{M} \langle dK, dK \rangle) (\mathbf{x}) \\ &\quad \times \left(n \frac{\partial B}{\partial \theta} + \frac{(n+1)^2}{8} \frac{\partial^2 B}{\partial s^2} + \frac{n^2-1}{8} \left(\frac{d_{1/2p}(K(\mathbf{x}))}{\sqrt{\varepsilon}} \right)^{-1} \frac{\partial B}{\partial s} \right) \\ &\quad + I_1^\varepsilon(\mathbf{x}) + I_2^\varepsilon(\mathbf{x}) + I_3^\varepsilon(\mathbf{x}) = \mathbb{G}_f^\varepsilon(\mathbf{x}) + \sum_{i=1}^4 I_i^\varepsilon(\mathbf{x}), \end{aligned} \quad (3.49)$$

where

$$I_1^\varepsilon(\mathbf{x}) = \frac{1}{2p} \varepsilon^{p-1/2} d_{1/2p-1} (K(\mathbf{x})) \left[\frac{\partial B}{\partial s} (\mathbb{M} (\mathcal{L}K)) (\mathbf{x}) + \frac{\partial^2 B}{\partial s \partial \theta} (\mathbb{M} \langle dK, d\Theta^\circ \rangle) (\mathbf{x}) \right],$$

$$I_2^\varepsilon(\mathbf{x}) = \varepsilon^p \left[\frac{\partial B}{\partial \theta} (\mathbb{M} (\mathcal{L}\Theta^\circ)) (\mathbf{x}) + \frac{1}{2} \frac{\partial^2 B}{\partial \theta^2} (\mathbb{M} \langle d\Theta^\circ, d\Theta^\circ \rangle) (\mathbf{x}) \right],$$

$$I_3^\varepsilon(\mathbf{x}) = n \varepsilon^{p-1} d_{n-1} (K(\mathbf{x})) (\mathbb{M} \langle dK, dK \rangle) (\mathbf{x}) \frac{\beta(0) - \beta(K(\mathbf{x}))}{\beta(K(\mathbf{x}))} \frac{\partial B}{\partial \theta},$$

$$I_4^\varepsilon(\mathbf{x}) = \varepsilon^{p-1} d_{n-1} (K(\mathbf{x})) \left(\frac{\partial \check{\Psi}_f}{\partial \theta} (\Theta(\mathbf{x}), K(\mathbf{x})) - \frac{\partial \check{\Psi}_f}{\partial \theta} (\Theta^\circ, 0) \right) (\mathbb{M} \langle dK, dK \rangle) (\mathbf{x}).$$

From this we see that

$$\lim_{\substack{\mathbf{y} \rightarrow \mathbf{x} \\ \mathbf{y} \in \mathcal{E} \setminus (\partial \mathbf{V} \cup \mathcal{C})}} (\mathbb{M} (\mathcal{L}^\varepsilon B^\varepsilon)) (\mathbf{y})$$

exists for all $\mathbf{x} \in \partial \mathbf{V} \cup \mathcal{C}$. Also, $(\mathbb{M} \langle dB^\varepsilon, dB^\varepsilon \rangle) (\mathbf{x}) = \sum_{i=5}^7 I_i^\varepsilon(\mathbf{x})$, where

$$I_5^\varepsilon(\mathbf{x}) = \varepsilon^{2p} \left(\frac{\partial B}{\partial \theta} \right)^2 (\mathbb{M} \langle d\Theta^\circ, d\Theta^\circ \rangle) (\mathbf{x}),$$

$$I_6^\varepsilon(\mathbf{x}) = \frac{1}{4p^2} \varepsilon^{2p-1} \left(\frac{\partial B}{\partial s} \right)^2 d_{n-1} (K(\mathbf{x})) (\mathbb{M} \langle dK, dK \rangle) (\mathbf{x}),$$

and

$$I_7^\varepsilon(\mathbf{x}) = \frac{1}{p} \varepsilon^{2p-1/2} \frac{\partial B}{\partial \theta} \frac{\partial B}{\partial s} \mathbf{d}_{1/2p-1}(K(\mathbf{x})) (\mathbb{M} \langle dK, d\Theta^\circ \rangle)(\mathbf{x}).$$

From Proposition 3.2, we know that $B \in C^2(\mathbb{R} \times [0, \infty))$ and

$$|\partial^\alpha B(\theta, s)| \leq C \left\| \check{\Psi}_f(\cdot, 0) \right\|_{C^5(\mathbb{R})} e^{-s/C} \quad (3.50)$$

for all $(\theta, s) \in \mathbb{R} \times (0, \infty)$, for some constant $C > 0$, and for any multi-index $\alpha, |\alpha| \leq 2$. Also, there exists a positive constant which does not depend of f such that

$$\left\| \check{\Psi}_f(\cdot, 0) \right\|_{C^5(\mathbb{R})} \leq C \|f\|_{C^5(\bar{\mathbf{I}})}. \quad (3.51)$$

Combining inequalities (3.50) and (3.51), we see that there exist constants $C_1, C_2 > 0$ such that

$$|\partial^\alpha B(\theta, s)| \leq C_1 \|f\|_{C^5(\bar{\mathbf{I}})} e^{-s/C_2} \quad (3.52)$$

for all $(\theta, s) \in \mathbb{R} \times (0, \infty)$, and for any multi-index $\alpha, |\alpha| \leq 2$. Now, for every $\mathbf{x} \in \mathcal{E} \setminus (\partial \mathbf{V} \cup \mathcal{C})$, there exist positive constants C_1, C_2 such that

$$|B^\varepsilon(\mathbf{x})| \leq \varepsilon^p |\partial^0 B(\theta, 0)| \leq C_1 \varepsilon^p \|f\|_{C^5(\bar{\mathbf{I}})},$$

and positive constants C_{ij} such that

$$\begin{aligned} |I_1^\varepsilon(\mathbf{x})| &\leq C_{11} \varepsilon^{p-1/2} \varepsilon^{p((1/2p)-1)} \|f\|_{C^5(\bar{\mathbf{I}})} \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\} \\ &= C_{12} \|f\|_{C^5(\bar{\mathbf{I}})} \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\}, \\ |I_2^\varepsilon(\mathbf{x})| &\leq C_{21} \varepsilon^p \|f\|_{C^5(\bar{\mathbf{I}})} \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\}, \\ |I_3^\varepsilon(\mathbf{x})| &\leq C_{31} n \varepsilon^{p-1} \left| \frac{\beta'(K(\mathbf{x}))}{\beta(K(\mathbf{x}))} \right| \mathbf{d}_n(K(\mathbf{x})) \|f\|_{C^5(\bar{\mathbf{I}})} \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\} \\ &\leq C_{32} n \varepsilon^{p-1+p} \|f\|_{C^5(\bar{\mathbf{I}})} \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\} \\ &\leq C_{33} \|f\|_{C^5(\bar{\mathbf{I}})} \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\}, \\ |I_4^\varepsilon(\mathbf{x})| &\leq C_{41} \varepsilon^p \|f\|_{C^5(\bar{\mathbf{I}})} \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\}, \\ |I_5^\varepsilon(\mathbf{x})| &\leq C_{51}^2 \varepsilon^{2p} \|f\|_{C^5(\bar{\mathbf{I}})}^2 \exp \left\{ -\frac{2|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\}, \\ |I_6^\varepsilon(\mathbf{x})| &\leq C_{61}^2 \varepsilon^{2p-1+p(n-1)} \|f\|_{C^5(\bar{\mathbf{I}})}^2 \exp \left\{ -\frac{2|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\} \\ &= C_{62}^2 \|f\|_{C^5(\bar{\mathbf{I}})}^2 \exp \left\{ -\frac{2|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\}, \end{aligned}$$

N. V. O'Bryant

and

$$|I_7^\varepsilon(\mathbf{x})| \leq C_{71}^2 \varepsilon^p \|f\|_{C^5(\bar{\mathbf{I}})}^2 \exp \left\{ -\frac{2|K(\mathbf{x})|^{1/2p}}{C\sqrt{\varepsilon}} \right\}.$$

Then, there exist positive constants C_1 and C_2 such that for every $\mathbf{x} \in \mathcal{E} \setminus (\partial \mathbf{V} \cup \mathcal{C})$

$$|(\mathbb{M}(\mathcal{L}^\varepsilon B^\varepsilon))(\mathbf{x}) - \mathbb{G}_f^\varepsilon(\mathbf{x})| \leq \sum_{i=1}^5 |I_i^\varepsilon(\mathbf{x})| \leq C_1 \|f\|_{C^5(\bar{\mathbf{I}})} \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C_2\sqrt{\varepsilon}} \right\}$$

and

$$\sqrt{(\mathbb{M}\langle dB^\varepsilon, dB^\varepsilon \rangle)(\mathbf{x})} \leq \sum_{i=5}^7 \sqrt{|I_i^\varepsilon(\mathbf{x})|} \leq C_1 \|f\|_{C^5(\bar{\mathbf{I}})} \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C_2\sqrt{\varepsilon}} \right\}.$$

This completes our proof of Lemma 3.6. $\square$

Examining expressions (3.43) and (3.49), we see that we would like to apply the martingale problem to a smoothed version of B^ε which we call Ψ^ε . From Proposition 3.6, B^ε is defined on $\mathcal{E}$ only, and its second derivatives are not guaranteed to be continuous on $\partial \mathbf{V} \cup \mathcal{C} \subset \mathcal{E}$. On the other hand, we would like to have $\Psi^\varepsilon \in C^2(\mathbb{R}^2)$.

First, we define $v_a \in C_c^\infty(\mathbb{R}; [0, 1])$ by $v_a(h) := v(3h/a)$, then

$$v_a(h) = \begin{cases} 1 & \text{if } |h| \leq a/3, \\ 0 & \text{if } |h| \geq 2a/3. \end{cases}$$

Using this function we cut off B^ε near the edges of $\mathcal{E}$ so that

$$\Psi^\varepsilon(\mathbf{x}) = v_a(K(\mathbf{x})) B^\varepsilon(\mathbf{x})$$

for every $\mathbf{x} \in \mathbb{R}^2$. Now, notice that

$$\begin{aligned} (\mathbb{M}(\mathcal{L}^\varepsilon \Psi^\varepsilon))(\mathbf{x}) - \mathbb{G}_f^\varepsilon(\mathbf{x}) &= v_a(K(\mathbf{x})) ((\mathbb{M}(\mathcal{L}^\varepsilon B^\varepsilon))(\mathbf{x}) - \mathbb{G}_f^\varepsilon(\mathbf{x})) \\ &\quad + B^\varepsilon(\mathbf{x}) \left(v_a'(K(\mathbf{x})) (\mathbb{M}(\mathcal{L}K))(\mathbf{x}) + \frac{1}{2} v_a''(K(\mathbf{x})) (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}) \right) \\ &\quad + v_a'(K(\mathbf{x})) (\mathbb{M}\langle dB^\varepsilon, dK \rangle)(\mathbf{x}) \end{aligned}$$

for all $\mathbf{x} \in \mathbb{R}^2 \setminus (\partial \mathbf{V} \cup \mathcal{C})$. Using Lemma 3.6, there exists a positive constant C such that

$$\begin{aligned} |(\mathbb{M}(\mathcal{L}^\varepsilon \Psi^\varepsilon))(\mathbf{x}) - \mathbb{G}_f^\varepsilon(\mathbf{x})| &\leq |((\mathbb{M}(\mathcal{L}^\varepsilon B^\varepsilon))(\mathbf{x}) - \mathbb{G}_f^\varepsilon(\mathbf{x}))| \\ &\quad + |B^\varepsilon(\mathbf{x})| \cdot \left| \beta'(K(\mathbf{x})) (\mathbb{M}(\mathcal{L}K))(\mathbf{x}) + \frac{1}{2} \beta''(K(\mathbf{x})) (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}) \right| \\ &\quad + |\beta'(K(\mathbf{x}))| \cdot |(\mathbb{M}\langle dB^\varepsilon, dK \rangle)(\mathbf{x})| \\ &\leq C \|f\|_{C^5(\bar{\mathbf{I}})} \left(\varepsilon^p + \exp \left\{ -\frac{|K(\mathbf{x})|^{1/2p}}{C_2\sqrt{\varepsilon}} \right\} \right). \end{aligned}$$

Notice that $\Psi^\varepsilon \in C^1(\mathbb{R}^2) \cup C^2(\mathbb{R}^2 \setminus (\partial\mathbf{V} \cup \mathcal{C}))$. Also,

$$\lim_{\substack{\mathbf{y} \rightarrow \mathbf{x} \\ \mathbf{y} \in \mathcal{E} \setminus (\partial\mathbf{V} \cup \mathcal{C})}} (\mathcal{L}_t^\varepsilon \Psi^\varepsilon)(\mathbf{y})$$

exists for all $t \geq 0$ and for all $\mathbf{x} \in \partial\mathbf{V} \cup \mathcal{C}$ since

$$\lim_{\substack{\mathbf{y} \rightarrow \mathbf{x} \\ \mathbf{y} \in \mathcal{E} \setminus (\partial\mathbf{V} \cup \mathcal{C})}} (\mathbb{M}(\mathcal{L}^\varepsilon \Psi^\varepsilon))(\mathbf{y})$$

exists for all $\mathbf{x} \in \partial\mathbf{V} \cup \mathcal{C}$.

We will use an even function $u_2 \in C_c^\infty(\mathbb{R}^2)$ with $\int_{\mathbb{R}^2} u_2(\mathbf{x}) d\mathbf{x} = 1$. For $\delta > 0$, we set the mollifier

$$\Psi_\delta^\varepsilon(\mathbf{y}) := \delta^{-2} \int_{\mathbb{R}^2} u_2\left(\frac{\mathbf{y} - \mathbf{z}}{\delta}\right) \Psi^\varepsilon(\mathbf{z}) d\mathbf{z}, \quad \mathbf{y} \in \mathbb{R}^2,$$

and observe that $\Psi_\delta^\varepsilon \in C^\infty(\mathbb{R}^2) \subset \mathcal{D}^\varepsilon$. Now, if $0 \leq t_1 < t_2 < \dots < t_m \leq s < t$, and $h_1, \dots, h_m \in C_b(\mathbb{R}^2)$, then

$$\mathbb{E}^\varepsilon \left[\left\{ (\Psi_\delta^\varepsilon)(\mathbf{X}_{t \wedge \tau}) - (\Psi_\delta^\varepsilon)(\mathbf{X}_{s \wedge \tau}) - \int_{s \wedge \tau}^{t \wedge \tau} \left(\mathcal{L}_{u/\varepsilon^2}^\varepsilon \Psi_\delta^\varepsilon \right)(\mathbf{X}_u) du \right\} \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] = 0.$$

Using properties of mollifiers, $\Psi_\delta^\varepsilon \rightarrow \Psi^\varepsilon$ a.e. as $\delta \rightarrow 0$, and

$$\mathcal{L}_t^\varepsilon \Psi_\delta^\varepsilon(\mathbf{y}) = \delta^{-2} \int_{\mathbb{R}^2} u_2\left(\frac{\mathbf{y} - \mathbf{z}}{\delta}\right) \mathcal{L}_t^\varepsilon \Psi^\varepsilon(\mathbf{z}) d\mathbf{z}.$$

Changing the order of integration,

$$\begin{aligned} & \mathbb{E}^\varepsilon \left[\left\{ \Psi^\varepsilon(\mathbf{X}_{t \wedge \tau}) - \Psi^\varepsilon(\mathbf{X}_{s \wedge \tau}) - \int_{s \wedge \tau}^{t \wedge \tau} \left(\mathcal{L}_{u/\varepsilon^2}^\varepsilon \Psi^\varepsilon \right)(\mathbf{X}_u) du \right\} \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \\ &= \lim_{\delta \rightarrow 0} \mathbb{E}^\varepsilon \left[\left\{ \Psi_\delta^\varepsilon(\mathbf{X}_{t \wedge \tau}) - \Psi_\delta^\varepsilon(\mathbf{X}_{s \wedge \tau}) - \int_{s \wedge \tau}^{t \wedge \tau} \left(\mathcal{L}_{u/\varepsilon^2}^\varepsilon \Psi_\delta^\varepsilon \right)(\mathbf{X}_u) du \right\} \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \\ &= 0. \end{aligned} \tag{3.53}$$

Applying Lemma 3.1, there exist positive constants C_1 and C_2 such that

$$\begin{aligned} & \left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} \left\{ \left(\mathcal{L}_{u/\varepsilon^2}^\varepsilon \Psi^\varepsilon \right)(\mathbf{X}_u) - (\mathbb{M}(\mathcal{L}^\varepsilon \Psi^\varepsilon))(\mathbf{X}_u) \right\} du \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right| \\ & < C_1 \varepsilon (1+t) \|\Psi^\varepsilon\|_{C^4(\bar{\mathbf{I}})} < C_2 \varepsilon^{p+2} \|f\|_{C^5(\bar{\mathbf{I}})}. \end{aligned}$$

Next, using the ideas described at the beginning of the proof of Proposition 3.3,

N. V. O'Bryant

and Eq. (3.53),

$$\begin{aligned}
& \left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} \left(f(\mathbf{X}_u) - (\mathbb{A}_0 f)([\mathbf{X}_u]) \right) d_n(K(\mathbf{X}_u)) \omega \left(\frac{K(\mathbf{X}_u)}{\varepsilon^p} \right) du \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right| \\
&= \left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} \varepsilon \mathbb{G}_f^\varepsilon(\mathbf{X}_u) du \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right| \\
&\leq \varepsilon \left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} \left(\mathbb{G}_f^\varepsilon(\mathbf{X}_u) - (\mathbb{M}(\mathcal{L}^\varepsilon \Psi^\varepsilon))(\mathbf{X}_u) \right) du \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right| \\
&+ \varepsilon \left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} \left((\mathbb{M}(\mathcal{L}^\varepsilon \Psi^\varepsilon))(\mathbf{X}_u) - (\mathcal{L}_{u/\varepsilon^2}^\varepsilon \Psi^\varepsilon)(\mathbf{X}_u) \right) du \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right| \\
&+ \varepsilon \left| \mathbb{E}^\varepsilon \left[\left(\Psi^\varepsilon(\mathbf{X}_{t \wedge \tau}) - \Psi^\varepsilon(\mathbf{X}_{s \wedge \tau}) \right) \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right|.
\end{aligned}$$

Therefore, there exists a positive constant C such that

$$\begin{aligned}
& \left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} \left(f(\mathbf{X}_u) - (\mathbb{A}_0 f)([\mathbf{X}_u]) \right) d_n(K(\mathbf{X}_u)) \omega \left(\frac{K(\mathbf{X}_u)}{\varepsilon^p} \right) du \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right| \\
&\leq C \varepsilon \|f\|_{C^5(\mathbf{\Gamma})} \left(\varepsilon^p + \varepsilon^{p+2} + \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} \left(\varepsilon^p + \exp \left\{ -\frac{|K(\mathbf{X}_u)|^{1/2p}}{C_2 \sqrt{\varepsilon}} \right\} \right) du \right] \right)
\end{aligned}$$

whenever $0 \leq t_1 < t_2 < \dots < t_m \leq s < t$, and $h_1, \dots, h_m \in C_b(\mathbb{R}^2)$. Finally, combining this inequality and inequality (3.36) with $v(h) = \exp \left\{ -\frac{1}{C_2} |h|^\alpha \right\}$, $\alpha = 1/2p$ and $\delta = \varepsilon^p$, we obtain (3.42). This completes the proof of Proposition 3.3. $\square$

4. The Martingale Problem

We want to prove that a family of probability measures converges to a unique solution $\mathbb{P}^*$ of the martingale problem for $(\mathcal{L}^*, \delta_{[\mathbf{x}]})$, and to identify the limiting generator $\mathcal{L}^*$ and its domain.

4.1. The limiting generator

Having the homeomorphism p into $\mathbb{R}^3$, we can define the metric ρ on $\mathbf{\Gamma}$ by

$$\rho([\mathbf{x}], [\mathbf{y}]) = \|p([\mathbf{x}]) - p([\mathbf{y}])\|_{\mathbb{R}^3}$$

for any $[\mathbf{x}], [\mathbf{y}] \in \mathbf{\Gamma}$. The constructed metric space $(\mathbf{\Gamma}, \rho)$ is Polish. Using Prohorov's theorem, we can show that the family of probability measures $\mathbb{P}^{\varepsilon,*} \in \mathcal{P}(C([0, \infty), \mathbf{\Gamma}))$ is tight in the Prohorov topology. We choose to omit the proof, since it is analogous to the proof of tightness in [?]. By Prohorov's theorem, tightness implies that $\{\mathbb{P}^{\varepsilon,*}\}$ has a cluster point. We will show that any such cluster point satisfies a certain martingale problem. Our next step is to identify the generator of the limiting martingale problem.

By properties of K , $K^{-1}(K(\mathbf{x})) = H^{-1}(H(\mathbf{x}))$ for any $\mathbf{x} \notin \overline{\mathbf{V}}$, and there exists a smooth extension of $K(\mathbf{x}) = (H(\mathbf{x}))^{1/n}$ inside $\overline{\mathbf{V}}$.

Given a function F and a set $[A] \in \mathbf{\Gamma}$, we denote the restriction of F to $[A]$ as F_A . In particular, if $F \in C(\mathbf{\Gamma}; \mathbb{R})$, then $F \circ \pi \in C(\overline{\mathbf{I}}, \mathbb{R})$ and

- $F_{\mathbf{V}} \circ \pi(\mathbf{x}) = F_{\mathbf{V}}(\mathbf{x})$ for $\mathbf{x} \in \mathbf{V}$;
- $F_{\overline{\mathbf{U}}} \circ \pi(\mathbf{x}) = (F_{\overline{\mathbf{U}}} \circ K^{-1}) \circ K$ for $\mathbf{x} \in \overline{\mathbf{U}}$;
- $F_K := F_{\overline{\mathbf{U}}} \circ K^{-1}$ maps $[0, K^*]$ into $\mathbb{R}$;
- $F \circ \pi(\mathbf{x}) = F_{\mathbf{V}}(\mathbf{x}) \mathbb{1}_{\mathbf{V}}(\mathbf{x}) + F_K(K(\mathbf{x})) \mathbb{1}_{\overline{\mathbf{I}} \setminus \mathbf{V}}(\mathbf{x})$ for $\mathbf{x} \in \overline{\mathbf{I}}$.

If $F \in C^2(\mathbf{\Gamma}_{\mathbf{V}} \cup \mathbf{\Gamma}_{\mathbf{U}}; \mathbb{R})$ then $F \circ \pi \in C^2(\mathbf{I} \setminus \partial \mathbf{V}; \mathbb{R})$, and for every $t \in \mathbb{R}$ the operator $\mathcal{L}_t(F \circ \pi)$ is well-defined on $\mathbf{I} \setminus \partial \mathbf{V}$. If $\mathbb{M}(\mathcal{L}(F \circ \pi)) \in \mathcal{D}_{\mathbb{A}}$, then for $[\mathbf{x}] \in \mathbf{\Gamma}_{\mathbf{V}} \cup \mathbf{\Gamma}_{\mathbf{U}}$, we set

$$(\mathbb{A}^* F)([\mathbf{x}]) = \begin{cases} \bar{b}(K(\mathbf{x})) F'_K(K(\mathbf{x})) + \frac{1}{2} \bar{\sigma}^2(K(\mathbf{x})) F''_K(K(\mathbf{x})) & \text{if } [\mathbf{x}] \in \mathbf{\Gamma}_{\mathbf{U}}, \\ \left(\mathbb{M}(\mathcal{L} F_{\mathbf{V}}) \right)(\mathbf{x}) & \text{if } [\mathbf{x}] \in \mathbf{\Gamma}_{\mathbf{V}}. \end{cases}$$

where

$$\bar{b} := \mathbb{A}_K \left(\mathbb{M} \left((\mathcal{L} K) - (\nabla \Phi_{\psi}, b) \right) \right)$$

and

$$\bar{\sigma}^2 := \mathbb{A}_K \left(\mathbb{M} \langle dK, dK \rangle \right).$$

Also, for future reference, define the second-order differential operator $\mathbb{M} \mathcal{L}^K$ on $C^2((0, K^*))$ by

$$(\mathbb{M}(\mathcal{L}^K f))(h) = \bar{b}(h) f'(h) + \frac{1}{2} \bar{\sigma}^2(h) f''(h).$$

For every $\mathbf{x} \in \mathbf{U} \setminus \overline{\mathbf{V}}$ and for every $F \in C^2(\mathbf{\Gamma}_{\mathbf{V}} \cup \mathbf{\Gamma}_{\mathbf{U}})$ such that $\mathbb{M}(\mathcal{L}(F \circ \pi)) \in \mathcal{D}_{\mathbb{A}}$, we have $(\mathbb{A}^* F)([\mathbf{x}]) = (\mathbb{M}(\mathcal{L}^K F_K))(K(\mathbf{x}))$.

We introduce two gluing operators. For F with $F_{\mathbf{V}} \in C^1(\mathbf{\Gamma}_{\mathbf{V}})$, we define an 'inner' gluing operator by

$$\begin{aligned} \underline{G}F &:= \lim_{\substack{[\mathbf{x}] \rightarrow \mathbf{\Gamma}_{\partial \mathbf{V}} \\ [\mathbf{x}] \in \mathbf{\Gamma}_{\mathbf{V}}}} \left(\mathbb{A}_0(\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right)([\mathbf{x}]) \\ &= \lim_{\substack{[\mathbf{x}] \rightarrow \mathbf{\Gamma}_{\partial \mathbf{V}} \\ [\mathbf{x}] \in \mathbf{\Gamma}_{\mathbf{V}}}} \left(\int_{\mathbf{y} \in [\mathbf{x}]} \frac{1}{\|\nabla K(\mathbf{y})\|} dl(\mathbf{y}) \right)^{-1} \int_{\mathbf{y} \in [\mathbf{x}]} \frac{(\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle)(\mathbf{y})}{\|\nabla K(\mathbf{y})\|} dl(\mathbf{y}), \end{aligned}$$

N. V. O'Bryant

if this limit exists. For F with $F_{\mathbf{U}} \in C^1(\Gamma_{\mathbf{U}})$, we define an 'outer' gluing operator

$$\begin{aligned}\overline{G}F &:= \lim_{\substack{[\mathbf{x}] \rightarrow \Gamma_{\partial\mathbf{V}} \\ [\mathbf{x}] \in \Gamma_{\mathbf{U}}}} \left(\mathbb{A}_0(\mathbb{M}\langle dF_K, dK \rangle) \right)([\mathbf{x}]) \\ &= \lim_{\substack{[\mathbf{x}] \rightarrow \Gamma_{\partial\mathbf{V}} \\ [\mathbf{x}] \in \Gamma_{\mathbf{U}}}} \left(\int_{\mathbf{y} \in [\mathbf{x}]} \frac{1}{\|\nabla K(\mathbf{y})\|} dl(\mathbf{y}) \right)^{-1} \int_{\mathbf{y} \in [\mathbf{x}]} \frac{(\mathbb{M}\langle dF_K, dK \rangle)(\mathbf{y})}{\|\nabla K(\mathbf{y})\|} dl(\mathbf{y}),\end{aligned}$$

if this limit exists.

Definition 4.2. Define the domain

$$\mathcal{D}^* := \left\{ F \in C(\Gamma) : F|_{\Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}}} \in C^2(\Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}}), \mathbb{M}(\mathcal{L}(F \circ \pi)) \in \mathcal{D}_{\mathbb{A}}, \right. \\ \left. \underline{G}F = \overline{G}F, \lim_{[\mathbf{y}] \rightarrow \Gamma_{\partial\mathbf{I}}} (\mathbb{A}^*F)([\mathbf{y}]) = 0 \right\}.$$

For any $F \in \mathcal{D}^*$, we define

$$(\mathcal{L}^*F)([\mathbf{x}]) := \lim_{\substack{[\mathbf{y}] \rightarrow [\mathbf{x}] \\ [\mathbf{y}] \in \Gamma_{\mathbf{U}} \cup \Gamma_{\mathbf{V}}}} (\mathbb{A}^*F)([\mathbf{y}]), \quad [\mathbf{x}] \in \Gamma.$$

4.2. Approximate test functions

If $F \in \mathcal{D}^*$, then $F \in C(\Gamma)$, and for all $\mathbf{x} \in \overline{\mathbf{I}}$,

$$F \circ \pi(\mathbf{x}) = F_{\mathbf{V}}(\mathbf{x}) \mathbb{1}_{\mathbf{V}}(\mathbf{x}) + F_K(K(\mathbf{x})) \mathbb{1}_{\overline{\mathbf{I}} \setminus \mathbf{V}}(\mathbf{x}).$$

Unfortunately, given $F \in \mathcal{D}^*$, we are not guaranteed that $F \circ \pi \in \mathcal{D}^{\varepsilon}$. According to the definition of $\mathcal{D}^*$, $F|_{\Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}}} \in C^2(\Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}})$, that is, function F has enough smoothness on both $\Gamma_{\mathbf{V}}$ and $\Gamma_{\mathbf{U}}$. But this does not immediately guarantee enough smoothness at $\Gamma_{\partial\mathbf{V}}$.

Proposition 4.4. *Let $F \in \mathcal{D}^*$, then $F_{\mathbf{V}} \in \mathbb{H}^2(\mathbf{V})$ and $F_K \in C^2([0, K^*])$.*

Proof. To study the regularity properties of F near $\Gamma_{\partial\mathbf{V}}$, we will use the continuity of $\mathcal{L}^*F$ on Γ .

Lemma 4.7. *Let $g \in C(\overline{\mathbf{V}})$ and $r \in \mathbb{R}$. There exists a unique solution $u_{g,r} \in \mathbb{H}^2(\mathbf{V}) \cap C(\overline{\mathbf{V}})$ of the PDE*

$$\mathbb{M}(\mathcal{L}u_{g,r}) = g \quad \text{in } \mathbf{V}, \quad u_{g,r} = r \quad \text{on } \partial\mathbf{V}. \quad (4.54)$$

If $g \in C^k(\overline{\mathbf{V}})$ and $k \geq 0$, then $u_{g,r} \in \mathbb{H}^{k+2}(\mathbf{V}) \cap C^k(\overline{\mathbf{V}})$ and there exists a constant C_k such that

$$\begin{aligned}\|u_{g,r}\|_{\mathbb{H}^{k+2}(\mathbf{V})} &\leq C_k \left\{ \|g\|_{C^k(\overline{\mathbf{V}})} + |r| \right\} \\ \|u_{g,r}\|_{C^k(\overline{\mathbf{V}})} &\leq C_k \left\{ \|g\|_{C^k(\overline{\mathbf{V}})} + |r| \right\}.\end{aligned} \quad (4.55)$$

Proof. Let $v \in \mathbb{H}_0^1(\mathbf{V})$ be a weak solution of the PDE

$$\begin{aligned} \mathbb{M}(\mathcal{L}v) &= 0 \quad \text{in } \mathbf{V}, \\ v &= 0 \quad \text{on } \partial\mathbf{V}. \end{aligned}$$

We know that $\partial\mathbf{V} \in C^\infty$ and all the coefficients of $\mathbb{M}\mathcal{L}$ are smooth. Then, by the infinite differentiability up to the boundary [?, Theorem 6, pg. 326], $v \in C^\infty(\overline{\mathbf{V}})$. By the weak maximum principle [?, Theorem 1, pg. 327], $v \equiv 0$ on $\overline{\mathbf{V}}$. The Fredholm alternative implies that a unique weak solution $u \in \mathbb{H}_0^1(\mathbf{V})$ of the corresponding inhomogeneous PDE (4.54) exists. By boundary $\mathbb{H}^2$ -regularity [?, Theorem 4, pg. 317], $u \in \mathbb{H}^2(\mathbf{V})$. By standard Sobolev theory, $u \in C(\overline{\mathbf{V}})$.

Now, we define $u_{g,r} := u + r$. Notice that this $u_{g,r}$ satisfies (4.54). Let $u'_{g,r}$ be another solution of (4.54), and take $u' = u_{g,r} - u'_{g,r}$ on $\overline{\mathbf{V}}$. Then $\mathbb{M}(\mathcal{L}u') = 0$ on $\mathbf{V}$, and by the infinite differentiability in the interior [?, Theorem 3, pg. 316], $u' \in C^\infty(\mathbf{V})$. On the other hand, $u'_{g,r} \in C(\overline{\mathbf{V}})$ implies that $u' \in C(\overline{\mathbf{V}})$, and by the maximum principle, $u' \equiv 0$ on $\overline{\mathbf{V}}$.

If $g \in C^k(\overline{\mathbf{V}})$, then $g \in \mathbb{H}^k(\mathbf{V})$, and by higher boundary regularity [?, Theorem 5, pg. 323], $u \in \mathbb{H}^{k+2}(\mathbf{V})$, and there exists a constant C_k such that inequalities (4.55) hold. $\square$

Lemma 4.8. Let $g \in C^1([0, K^*])$ and $r_1, r_2 \in \mathbb{R}$.

There exists a unique solution $u_{g,r_1,r_2} \in C^2([0, K^*])$ of the PDE

$$\mathbb{M}(\mathcal{L}^K u_{g,r_1,r_2}) = g \text{ on } (0, K^*), \quad u_{g,r_1,r_2}(0) = r_1, \quad u_{g,r_1,r_2}(K^*) = r_2. \quad (4.56)$$

If $g \in C^k([0, K^*])$ and $k \geq 0$, then $u_{g,r_1,r_2} \in C^{k+1}([0, K^*])$ and there exists a constant C_k such that

$$\|u_{g,r_1,r_2}\|_{C^{k+1}([0, K^*])} \leq C_k \left\{ \|g\|_{C^k([0, K^*])} + |r_1| + |r_2| \right\}. \quad (4.57)$$

Proof. Let

$$I(h) = \exp \left\{ 2 \int_0^h \frac{\overline{b}(s)}{\overline{\sigma}^2(s)} ds \right\}, \quad h \in [0, K^*].$$

Notice that

$$\begin{aligned} u_{g,r_1,r_2}(h) &= \frac{1}{\int_0^{K^*} \frac{ds}{I(s)}} \left(r_1 \int_h^{K^*} \frac{ds}{I(s)} + r_2 \int_0^h \frac{ds}{I(s)} \right) \\ &\quad + 2 \int_0^h \frac{1}{I(q)} \left(\int_0^q \frac{g(s)}{\overline{\sigma}^2(s)} I(s) ds \right) dq \\ &\quad - 2 \frac{\int_0^h \frac{ds}{I(s)}}{\int_0^{K^*} \frac{ds}{I(s)}} \int_0^{K^*} \frac{1}{I(q)} \left(\int_0^q \frac{g(s)}{\overline{\sigma}^2(s)} I(s) ds \right) dq, \end{aligned}$$

solves the PDE (4.56) uniquely.

If $g \in C^k([0, K^*])$, then $u_{g,r_1,r_2} \in C^{k+1}([0, K^*])$ and there exists a constant C_k such that (4.57) holds. $\square$

N. V. O'Bryant

Now, apply Lemma 4.7 with $g = \mathbb{M}(\mathcal{L}F_{\mathbf{V}})$ and $r = F(\partial\mathbf{V})$ to see that $u_{g,r} = F_{\mathbf{V}} \in \mathbb{H}^2(\mathbf{V})$. Lemma 4.8 with $g = \mathbb{M}(\mathcal{L}^K F_K)$, $r_1 = F_K(0)$, and $r_2 = F_K(K^*)$ implies that $u_{g,r_1,r_2} = F_K \in C^2([0, K^*])$. Therefore, the proof of Proposition 4.4 is complete. $\square$

Although $F \circ \pi$ itself is not guaranteed to belong to $\mathcal{D}^\varepsilon$, it can be approximated by a collection of functions $\{F^\varepsilon, \varepsilon > 0\}$ from $C^2(\mathbb{R}^2) \subset \mathcal{D}^\varepsilon$, so that

$$\lim_{\varepsilon \rightarrow 0} \sup_{\mathbf{x} \in \bar{\mathbf{I}}} |F^\varepsilon(\mathbf{x}) - F \circ \pi(\mathbf{x})| = 0.$$

We will define F^ε similarly to the approximate test functions defined in [?]. The special construction of these functions is sketched below and will be used later.

First, we define the cutoff function $v_+ \in C^\infty(\mathbb{R})$ by

$$v_+(h) := \begin{cases} 0 & \text{if } h \leq 1, \\ 1 & \text{if } h \geq 2. \end{cases}$$

Next, we build suitable extensions of the functions $F_{\mathbf{V}}$ and F_K .

Lemma 4.9. *Let $F \in \mathcal{D}^*$. Suppose that $\partial\mathbf{V} \subset Z_1 \subset \mathbb{R}^2$. There exists a family of functions $\{F_{\mathbf{V}}^\varepsilon, \varepsilon > 0\}$, such that every $F_{\mathbf{V}}^\varepsilon \in C_c^8(Z_1)$, and*

$$\begin{aligned} F_{\mathbf{V}}^\varepsilon(\mathbf{x}) &= F(\mathbf{T}_{\partial\mathbf{V}}) \text{ for all } \mathbf{x} \in \partial\mathbf{V}, \varepsilon > 0, \\ \lim_{\varepsilon \rightarrow 0} \|F_{\mathbf{V}}^\varepsilon - F_{\mathbf{V}}\|_{\mathbb{H}^2(\mathbf{V})} &= 0, \\ \lim_{\varepsilon \rightarrow 0} \|\mathbb{M}(\mathcal{L}F_{\mathbf{V}}^\varepsilon) - \mathcal{L}^*F_{\mathbf{V}}\|_{C(\bar{\mathbf{V}})} &= 0, \\ \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon^{kq} \|F_{\mathbf{V}}^\varepsilon\|_{C^k(Z_1)} &< \infty, k \leq 8. \end{aligned}$$

Lemma 4.10. *Let $F \in \mathcal{D}^*$. Suppose that $[0, K^*] \subset Z_2 \subset \mathbb{R}$. There exists a sequence of functions $\{F_K^\varepsilon, \varepsilon > 0\}$ such that every $F_K^\varepsilon \in C_c^8(Z_2)$ and*

$$\begin{aligned} F_K^\varepsilon(0) &= F(\mathbf{T}_{\partial\mathbf{V}}) \text{ for all } \varepsilon > 0, \\ \lim_{\varepsilon \rightarrow 0} \|F_K^\varepsilon - F_K\|_{C^2([0, K^*])} &= 0, \\ \lim_{\varepsilon \rightarrow 0} \left\| \mathbb{M}(\mathcal{L}^K F_K^\varepsilon) - (\mathcal{L}^*F_K)^K \right\|_{C([0, K^*])} &= 0, \\ \overline{\lim}_{\varepsilon \rightarrow 0} \varepsilon^{kq} \|F_K^\varepsilon\|_{C^k(Z_2)} &< \infty, k \leq 8. \end{aligned}$$

Lemma 4.9 and Lemma 4.10 can be proved in the same fashion as Lemma 4.5 and Lemma 4.6 from [?], but instead of the properties of the generator $\mathcal{L}$ from [?], we have to pay attention to the continuity properties of the time average of $\mathcal{L}_{t/\varepsilon}^\varepsilon$.

Finally, given any $\mathbf{x} \in \bar{\mathbf{I}}, \varepsilon > 0$, we let

$$F^\varepsilon(\mathbf{x}) := F_{\mathbf{V}}^\varepsilon(\mathbf{x}) \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \right) + F_K^\varepsilon(K(\mathbf{x})) v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right), \quad (4.58)$$

where $F_{\mathbf{V}}^\varepsilon$ is constructed in Lemma 4.9, and F_K^ε is constructed in Lemma 4.10.

4.3. Gluing

The hypothesis that $\underline{G}F = \overline{G}F$ will be called the ‘gluing requirement’. It can be rewritten in the form

$$\lim_{\substack{[\mathbf{x}] \rightarrow \Gamma_{\partial \mathbf{V}} \\ [\mathbf{x}] \in \Gamma_{\mathbf{V}}}} \left(\mathbb{A}_0 (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) ([\mathbf{x}]) = \lim_{\substack{[\mathbf{x}] \rightarrow \Gamma_{\partial \mathbf{V}} \\ [\mathbf{x}] \in \Gamma_{\mathbf{U}}}} \left(\mathbb{A}_0 (\mathbb{M} \langle dF_K, dK \rangle) \right) ([\mathbf{x}]).$$

Notice that if $[\mathbf{x}] \in \Gamma_{\mathbf{V}}$, then

$$\left(\mathbb{A}_0 (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) ([\mathbf{x}]) = \left(\mathbb{A}_K (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (K(\mathbf{x})).$$

If $[\mathbf{x}] \in \Gamma_{\mathbf{U}}$, then

$$\begin{aligned} \left(\mathbb{A}_0 (\mathbb{M} \langle dF_K, dK \rangle) \right) ([\mathbf{x}]) &= F'_K(K(x)) \left(\mathbb{A}_0 (\mathbb{M} \langle dK, dK \rangle) \right) ([\mathbf{x}]) \\ &= \overline{\sigma}^2(K(\mathbf{x})) F'_K(K(\mathbf{x})). \end{aligned}$$

So, formally, the gluing requirement demands that

$$\lim_{h \uparrow 0} \left(\mathbb{A}_K (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (h) = \lim_{h \downarrow 0} \overline{\sigma}^2(h) F'_K(h).$$

Proposition 4.5. *Let $F \in C(\Gamma)$ be such that $F|_{\Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}}} \in C^2(\Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}})$ and $\mathbb{M}(\mathcal{L}(F \circ \pi)) \in \mathcal{D}_{\mathbb{A}}$, then both $\underline{G}F$ and $\overline{G}F$ exist.*

Proof. From Lemma 4.4, $\lim_{h \downarrow 0} (F_K)'(h)$ exists; we denote this limit by $(F_K)'(0)$. Notice that from our assumptions on function K the limit $\lim_{h \downarrow 0} \overline{\sigma}^2(h) = \overline{\sigma}^2(0)$ exists and $\overline{\sigma}^2(0) > 0$. Then the quantity $\overline{G}F$ is well-defined and

$$\overline{G}F = \overline{\sigma}^2(0) (F_K)'(0).$$

For every $h \in (-a, a)$, we introduce a bounded linear operator $\mathbf{T}_h : \mathbb{H}^1(\mathbf{V}) \rightarrow L^2(K^{-1}(h))$ such that if $f \in \mathbb{H}^1(\mathbf{V}) \cap C(\overline{\mathbf{V}})$ then $\mathbf{T}_h f = f|_{\partial \mathbf{V}}$ and

$$\|\mathbf{T}_h f\|_{L^2(\partial \mathbf{V})} \leq C \|f\|_{\mathbb{H}^1(\mathbf{V})}. \quad (4.59)$$

The Trace theorem [?, Theorem 1, pg. 258] guarantees the existence of such operators $\mathbf{T}_h$. For any $f \in \mathbb{H}^1(\mathbf{V})$ and for $h \in (-a, 0]$ we define

$$(\mathbb{A}^{\text{Tr}} f)(h) = \left(\int_{\mathbf{y} \in K^{-1}(h)} \frac{1}{\|\nabla K(\mathbf{y})\|} dl(\mathbf{y}) \right)^{-1} \int_{\mathbf{y} \in K^{-1}(h)} \frac{(\mathbf{T}_h f)(\mathbf{y})}{\|\nabla K(\mathbf{y})\|} dl(\mathbf{y}).$$

If $f \in C(\overline{\mathbf{V}})$, then $\mathbb{A}^{\text{Tr}} f = A_K f$ on $(-a, 0]$. By Lemma 4.4, $F_{\mathbf{V}} \in \mathbb{H}^2(\mathbf{V})$, which implies that $\left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (0)$ is well-defined and that

$$\underline{G}F = \lim_{h \uparrow 0} \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (h), \quad (4.60)$$

N. V. O'Bryant

if this limit exists. We consider the sequence of functions $F_{\mathbf{V}}^\varepsilon$ from Lemma 4.9. By property (4.59) of the trace operator, there exists a constant $C > 0$ such that

$$\left| \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (h) - \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle) \right) (h) \right| \leq C \|F_{\mathbf{V}} - F_{\mathbf{V}}^\varepsilon\|_{\mathbb{H}^2(\mathbf{V})}$$

for all $h \in (-a, 0]$.

The following result is borrowed from [?] (Lemma 9.6).

Lemma 4.11. *There exists a constant $C > 0$ such that if $f \in \mathbb{H}^2(\mathbb{R}^2)$, then*

$$\left| \left(\mathbb{A}^{\text{Tr}} f \right) (h') - \left(\mathbb{A}^{\text{Tr}} f \right) (h'') \right| \leq C \|f\|_{\mathbb{H}^1(\mathbb{R}^2)} \sqrt{|h' - h''|} \quad (4.61)$$

for all $h', h'' \in (-a, a)$.

From (4.61), there exists a constant $C > 0$ such that

$$\begin{aligned} & \left| \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (h) - \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (0) \right| \\ & \leq \left| \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (h) - \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle) \right) (h) \right| \\ & \quad + \left| \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle) \right) (h) - \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle) \right) (0) \right| \\ & \quad + \left| \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle) \right) (0) - \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (0) \right| \\ & \leq 2C \|F_{\mathbf{V}} - F_{\mathbf{V}}^\varepsilon\|_{\mathbb{H}^2(\mathbf{V})} + C \sqrt{|h|} \|F_{\mathbf{V}}^\varepsilon\|_{\mathbb{H}^2(\mathbf{V})}. \end{aligned}$$

Therefore the limit in (4.60) exists and $\underline{G}F = \left(\mathbb{A}^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right) (0)$ is well-defined. $\square$

4.4. The limiting martingale problem

To prove that any cluster point of the $\mathbb{P}^{\varepsilon,*}$ s satisfies the martingale problem for $(\mathcal{L}^*, \delta_{[\mathbf{x}]})$, we will need to connect elements of the limiting domain $\mathcal{D}^*$ back to the functions from $\mathcal{D}^\varepsilon$. The propositions and lemmas below allow us to establish this connections.

Recall F^ε , constructed above to satisfy (4.58).

Proposition 4.6. *Given $\mathbf{x} \in \bar{\mathbf{I}}$, for every $t \geq 0$ and for every $\varepsilon > 0$*

$$\left(\mathcal{L}_{t/\varepsilon^2}^\varepsilon F^\varepsilon \right) (\mathbf{x}) = L^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) + \sum_{k=0}^3 G_k^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) + R^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right), \quad (4.62)$$

where

$$\begin{aligned}
L^\varepsilon(\mathbf{x}, t) &= \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right)\right) (\mathcal{L}_t F_{\mathbf{V}}^\varepsilon)(\mathbf{x}) + v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \mathcal{L}_t F_K^\varepsilon(K(\mathbf{x})) \\
&\quad + \frac{1}{\varepsilon} v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) (F_K^\varepsilon)'(K(\mathbf{x})) (\nabla K(\mathbf{x}), b(\mathbf{x}, t)), \\
G_0^\varepsilon(\mathbf{x}, t) &= \frac{1}{\varepsilon^{p+1}} v_+' \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) (\nabla K(\mathbf{x}), b(\mathbf{x}, t)) \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})\right), \\
G_1^\varepsilon(\mathbf{x}, t) &= \frac{1}{\varepsilon^p} v_+' \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \left(\langle dF_{\mathbf{V}}^\varepsilon, dK \rangle_t(\mathbf{x}) - \langle d(F_K^\varepsilon \circ K), dK \rangle_t(\mathbf{x})\right), \\
G_2^\varepsilon(\mathbf{x}, t) &= \frac{1}{2\varepsilon^{2p}} v_+'' \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})\right) \langle dK, dK \rangle_t(\mathbf{x}), \\
G_3^\varepsilon(\mathbf{x}, t) &= \frac{1}{\varepsilon} \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right)\right) (\nabla F_{\mathbf{V}}^\varepsilon(\mathbf{x}), b(\mathbf{x}, t)), \\
R^\varepsilon(\mathbf{x}, t) &= \frac{1}{\varepsilon^p} v_+' \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})\right) (\mathcal{L}_t K)(\mathbf{x}).
\end{aligned}$$

Proof. Below, for any appropriate f and g involved, we understand $\mathcal{L}_{t/\varepsilon^2}^\varepsilon f(g(\mathbf{x}))$ and $df(g(\mathbf{x}))$ as $(\mathcal{L}_{t/\varepsilon^2}^\varepsilon(f \circ g))(\mathbf{x})$ and $d(f \circ g)(\mathbf{x})$ correspondingly. For example,

$$\begin{aligned}
\mathcal{L}_{t/\varepsilon^2}^\varepsilon v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) &= \frac{1}{\varepsilon} \left(\nabla v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right), b\left(\mathbf{x}, \frac{t}{\varepsilon^2}\right)\right) + \mathcal{L}_{t/\varepsilon^2} v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \\
&= \frac{1}{\varepsilon^{p+1}} v_+' \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \left(\nabla K(\mathbf{x}), b\left(\mathbf{x}, \frac{t}{\varepsilon^2}\right)\right) \\
&\quad + \frac{1}{2} \sum_{i,j=1}^2 \gamma_{ij} \left(\mathbf{x}, \frac{t}{\varepsilon^2}\right) \frac{\partial^2 v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right)}{\partial x_i \partial x_j}(\mathbf{x}).
\end{aligned}$$

This implies that

$$\begin{aligned}
\mathcal{L}_{t/\varepsilon^2}^\varepsilon v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) &= \frac{1}{\varepsilon^{p+1}} v_+' \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \left(\nabla K(\mathbf{x}), b\left(\mathbf{x}, \frac{t}{\varepsilon^2}\right)\right) \\
&\quad + \frac{1}{\varepsilon^p} v_+' \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) (\mathcal{L}_{t/\varepsilon^2} K)(\mathbf{x}) \\
&\quad + \frac{1}{2\varepsilon^{2p}} v_+'' \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \langle dK, dK \rangle_{t/\varepsilon^2}(\mathbf{x}) \tag{4.63}
\end{aligned}$$

and

$$dv_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) = \frac{1}{\varepsilon^p} v_+' \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) dK(\mathbf{x}). \tag{4.64}$$

We know that $F^\varepsilon \in C^2(\mathbb{R}^2) \subset \mathcal{D}^\varepsilon$. Applying the generator $\mathcal{L}_{t/\varepsilon^2}^\varepsilon$ to the function

N. V. O'Bryant

F^ε defined in (4.58), we obtain

$$\begin{aligned}
(\mathcal{L}_{t/\varepsilon^2}^\varepsilon F^\varepsilon)(\mathbf{x}) &= \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right)\right) \left(\frac{1}{\varepsilon} \left(\nabla F_{\mathbf{V}}^\varepsilon(\mathbf{x}), b\left(\mathbf{x}, \frac{t}{\varepsilon^2}\right)\right) + (\mathcal{L}_{t/\varepsilon^2}^\varepsilon F_{\mathbf{V}}^\varepsilon)(\mathbf{x})\right) \\
&\quad + v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \mathcal{L}_{t/\varepsilon^2}^\varepsilon F_K^\varepsilon(K(\mathbf{x})) \\
&\quad + \frac{1}{\varepsilon} v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) (F_K^\varepsilon)'(K(\mathbf{x})) \left(\nabla K(\mathbf{x}), b\left(\mathbf{x}, \frac{t}{\varepsilon^2}\right)\right) \\
&\quad + \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})\right) \mathcal{L}_{t/\varepsilon^2}^\varepsilon v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \\
&\quad - \left\langle dF_{\mathbf{V}}^\varepsilon(\mathbf{x}), dv_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \right\rangle_{t/\varepsilon^2} \\
&\quad + \left\langle dF_K^\varepsilon(K(\mathbf{x})), dv_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \right\rangle_{t/\varepsilon^2}.
\end{aligned}$$

We unravel this expression further, using (4.63) and (4.64), to see that

$$\begin{aligned}
(\mathcal{L}_{t/\varepsilon^2}^\varepsilon F^\varepsilon)(\mathbf{x}) &= \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right)\right) \left(\frac{1}{\varepsilon} \left(\nabla F_{\mathbf{V}}^\varepsilon(\mathbf{x}), b\left(\mathbf{x}, \frac{t}{\varepsilon^2}\right)\right) + (\mathcal{L}_{t/\varepsilon^2}^\varepsilon F_{\mathbf{V}}^\varepsilon)(\mathbf{x})\right) \\
&\quad + v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \mathcal{L}_{t/\varepsilon^2}^\varepsilon F_K^\varepsilon(K(\mathbf{x})) \\
&\quad + \frac{1}{\varepsilon} v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) (F_K^\varepsilon)'(K(\mathbf{x})) \left(\nabla K(\mathbf{x}), b\left(\mathbf{x}, \frac{t}{\varepsilon^2}\right)\right) \\
&\quad + \frac{1}{\varepsilon^{p+1}} v'_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \left(\nabla K(\mathbf{x}), b\left(\mathbf{x}, \frac{t}{\varepsilon^2}\right)\right) \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})\right) \\
&\quad + \frac{1}{\varepsilon^p} v'_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})\right) (\mathcal{L}_{t/\varepsilon^2}^\varepsilon K)(\mathbf{x}) \\
&\quad + \frac{1}{2\varepsilon^{2p}} v''_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})\right) \langle dK, dK \rangle_{t/\varepsilon^2}(\mathbf{x}) \\
&\quad - \frac{1}{\varepsilon^p} v'_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) \left(\langle dF_{\mathbf{V}}^\varepsilon, dK \rangle_{t/\varepsilon^2}(\mathbf{x}) - \langle d(F_K^\varepsilon \circ K), dK \rangle_{t/\varepsilon^2}(\mathbf{x})\right).
\end{aligned}$$

This immediately implies (4.62). $\square$

Proposition 4.7. *Fix $F \in \mathcal{D}^*$ and $t > 0$. Then*

$$\overline{\lim}_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((\mathbb{M}L^\varepsilon)(\mathbf{X}_s) - (\mathcal{L}^*F)([\mathbf{X}_s]) \right) ds \right| \right] = 0. \quad (4.65)$$

Proof. Consider

$$\begin{aligned}
(\mathbb{M}L^\varepsilon)(\mathbf{x}) &= \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right)\right) (\mathbb{M}(\mathcal{L}F_{\mathbf{V}}^\varepsilon))(\mathbf{x}) \\
&\quad + v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) (\mathbb{M}(\mathcal{L}(F_K^\varepsilon \circ K)))(\mathbf{x}) \\
&= \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right)\right) (\mathbb{M}(\mathcal{L}F_{\mathbf{V}}^\varepsilon))(\mathbf{x}) \\
&\quad + v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) (F_K^\varepsilon)'(K(\mathbf{x})) (\mathbb{M}(\mathcal{L}K))(\mathbf{x}) \\
&\quad + \frac{1}{2} v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right) (F_K^\varepsilon)''(K(\mathbf{x})) (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}).
\end{aligned}$$

Recall that $v, v_+ \in C^\infty(\mathbb{R}; [0, 1])$ were defined to satisfy

$$v(h) = \begin{cases} 0 & \text{if } |h| \leq 1, \\ 1 & \text{if } |h| \geq 2 \end{cases} \quad \text{and} \quad v_+(h) = \begin{cases} 0 & \text{if } h \leq 1, \\ 1 & \text{if } h \geq 2. \end{cases}$$

Define $v_- \in C^\infty(\mathbb{R}; [0, 1])$ by $v_-(h) := v_+(-h)$, then

$$v_-(h) = \begin{cases} 1 & \text{if } h \leq -2, \\ 0 & \text{if } |h| \geq -1. \end{cases}$$

Without loss of generality assume that these three functions v, v_- and v_+ form a smooth partition of unity. Notice that

$$\text{supp}(v_-) \subset (-\infty, -1), \text{supp}(v) \subset (-2, 2), \text{supp}(v_+) \subset (1, \infty).$$

Fix r so that $\varepsilon^{2q} \gg \varepsilon^r \gg \varepsilon^p$ as $\varepsilon \rightarrow 0$. Then we have $\text{supp}\left(v_- \left(\frac{K(\cdot)}{\varepsilon^r}\right)\right) \subseteq \mathbf{V}$ and $\text{supp}\left(v_+ \left(\frac{K(\cdot)}{\varepsilon^r}\right)\right) \subseteq \mathbf{U}$. Notice that

$$v_- \left(\frac{K(\mathbf{x})}{\varepsilon^r}\right) \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p}\right)\right) = v_- \left(\frac{K(\mathbf{x})}{\varepsilon^r}\right),$$

which implies that

$$(\mathbb{M}L^\varepsilon)(\mathbf{x}) v_- \left(\frac{K(\mathbf{x})}{\varepsilon^r}\right) = (\mathbb{M}(\mathcal{L}F_{\mathbf{V}}^\varepsilon))(\mathbf{x}) v_- \left(\frac{K(\mathbf{x})}{\varepsilon^r}\right). \quad (4.66)$$

Now, using Lemma 4.9 and expression (4.66)

$$\begin{aligned}
\overline{\lim}_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((\mathbb{M}L^\varepsilon)(\mathbf{X}_u) - (\mathcal{L}^*F)(\mathbf{Y}_u) \right) v_- \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r}\right) du \right| \right] \\
\leq \overline{\lim}_{\varepsilon \rightarrow 0} \left\| \mathbb{M}(\mathcal{L}F_{\mathbf{V}}^\varepsilon) - \mathcal{L}^*F \right\|_{C(\overline{\mathbf{V}})} = 0. \quad (4.67)
\end{aligned}$$

N. V. O'Bryant

Next, by Lemma 4.9, Lemma 4.10, and using the fact that $\mathcal{L}^*F$ is bounded, we see that

$$\left| \left((\mathbb{M}L^\varepsilon)(\mathbf{x}) - (\mathcal{L}^*F)([\mathbf{x}]) \right) v \left(\frac{K(\mathbf{x})}{\varepsilon^r} \right) \right| < C\varepsilon^{-2q} \chi_{[-2\varepsilon^r, 2\varepsilon^r]}(K(\mathbf{x})).$$

By Lemma 3.5, there exists a positive constant C such that

$$\begin{aligned} \overline{\lim}_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((\mathbb{M}L^\varepsilon)(\mathbf{X}_u) - (\mathcal{L}^*F)(\mathbf{Y}_u) \right) v \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] \\ \leq C\varepsilon^{r-2q} \overline{\lim}_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\frac{1}{\varepsilon^r} \int_0^{t \wedge \tau} \chi_{[-2\varepsilon^r, 2\varepsilon^r]}(K(\mathbf{X}_u)) du \right] = 0. \end{aligned} \quad (4.68)$$

Notice that

$$\begin{aligned} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((\mathbb{M}L^\varepsilon)(\mathbf{X}_u) - (\mathcal{L}^*F)(\mathbf{Y}_u) \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] \\ \leq \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((\mathbb{M}L^\varepsilon)(\mathbf{X}_u) - (A_0(\mathbb{M}L^\varepsilon))(\mathbf{Y}_u) \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] \\ + \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((A_0(\mathbb{M}L^\varepsilon))(\mathbf{Y}_u) - L^\varepsilon \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] \\ + \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left(L^\varepsilon \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) - (\mathcal{L}^*F)(\mathbf{Y}_u) \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right], \end{aligned} \quad (4.69)$$

where

$$\begin{aligned} & \left(L^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) - (\mathcal{L}^*F)([\mathbf{x}]) \right) v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^r} \right) \\ &= \left(\mathcal{L}_{t/\varepsilon^2} F_K^\varepsilon(K(\mathbf{x})) + \frac{1}{\varepsilon} (F_K^\varepsilon)'(K(\mathbf{x})) \psi \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) \right) v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^r} \right) \\ &\quad - \mathbb{A}_0 \left(\mathbb{M}((\mathcal{L}K) - (\nabla \Phi_\psi, b)) \right) ([\mathbf{x}]) F_K'(K(\mathbf{x})) v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^r} \right) \\ &\quad - \frac{1}{2} \mathbb{A}_0 \left(\mathbb{M}(\langle dK, dK \rangle) \right) ([\mathbf{x}]) F_K''(K(\mathbf{x})) v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^r} \right) \\ &= v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^r} \right) \sum_{i=1}^5 K_i^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) \end{aligned} \quad (4.70)$$

and

$$\begin{aligned}
K_1^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) &= \left(\mathcal{L}_{t/\varepsilon^2} F_K^\varepsilon (K(\mathbf{x})) - (\mathbb{A}_0 (\mathbb{M} (\mathcal{L} (F_K^\varepsilon \circ K))) ([\mathbf{x}])) \right), \\
K_2^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) &= \left((\mathbb{A}_0 (\mathbb{M} (\mathcal{L} (F_K^\varepsilon \circ K))) ([\mathbf{x}])) - (\mathbb{A}_0 (\mathbb{M} (\mathcal{L} (F_K \circ K))) ([\mathbf{x}])) \right), \\
K_3^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) &= \left(F_K' (K(\mathbf{x})) - (F_K^\varepsilon)' (K(\mathbf{x})) \right) \mathbb{A}_0 (\mathbb{M} (\nabla \Phi_\psi, b)) ([\mathbf{x}]), \\
K_4^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) &= \left(\mathbb{A}_0 (\mathbb{M} (\nabla \Phi_\psi, b)) ([\mathbf{x}])) - (\nabla \Phi_\psi, b) \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) \right) (F_K^\varepsilon)' (K(\mathbf{x})), \\
K_5^\varepsilon \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) &= \left(\frac{1}{\varepsilon} \psi \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) + (\nabla \Phi_\psi, b) \left(\mathbf{x}, \frac{t}{\varepsilon^2} \right) \right) (F_K^\varepsilon)' (K(\mathbf{x})).
\end{aligned}$$

By Lemma 3.3, there exists a constant $C_c > 0$ such that

$$\begin{aligned}
\mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} K_1^\varepsilon \left(\mathbf{X}_u, \frac{t}{\varepsilon^2} \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] \\
\leq C_c \varepsilon^{1-r(n+1)} (1+t) \|F_K^\varepsilon\|_{C^5([0, K^*])}. \quad (4.71)
\end{aligned}$$

By Lemma 4.10,

$$\overline{\lim}_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} K_i^\varepsilon \left(\mathbf{X}_u, \frac{t}{\varepsilon^2} \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] = 0, \quad i = 2, 3. \quad (4.72)$$

By Lemma 3.3 and Lemma 4.10, there exists a constant $C > 0$ such that

$$\begin{aligned}
\mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} K_4^\varepsilon \left(\mathbf{X}_u, \frac{t}{\varepsilon^2} \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] \\
< C \varepsilon^{1-r(n+1)} (1+t) \|F_K^\varepsilon\|_{C^1([0, K^*])}. \quad (4.73)
\end{aligned}$$

By Lemma 3.4, there exists a constant $C_b > 0$ such that

$$\begin{aligned}
\mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} K_5^\varepsilon \left(\mathbf{X}_u, \frac{t}{\varepsilon^2} \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] \\
< C_b \varepsilon^{1-2r} (1+t) \|F_K^\varepsilon\|_{C^3([0, K^*])}. \quad (4.74)
\end{aligned}$$

Combining equality (4.70) with inequalities (4.71), (4.72), (4.73), and (4.74), we can see that

$$r < \frac{1-4q}{n+1}$$

guarantees that

$$\overline{\lim}_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left(L^\varepsilon \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) - (\mathcal{L}^* F) ([\mathbf{X}_u]) \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] = 0. \quad (4.75)$$

N. V. O'Bryant

By Lemma 3.2, Lemma 3.3, and Lemma 4.10, there exists a positive constant C such that

$$\begin{aligned} & \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((\mathbb{M}L^\varepsilon)(\mathbf{X}_u) - (A_0(\mathbb{M}L^\varepsilon))([\mathbf{X}_u]) \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] \\ & + \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((A_0(\mathbb{M}L^\varepsilon))([\mathbf{X}_u]) - L^\varepsilon \left(\mathbf{X}_u, \frac{u}{\varepsilon^2} \right) \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] \\ & < C\varepsilon^{1-r(n+1)-5q}(1+t) \end{aligned} \quad (4.76)$$

Therefore, combining (4.69), (4.75), and (4.76), we conclude that

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} \left((\mathbb{M}L^\varepsilon)(\mathbf{X}_u) - (\mathcal{L}^*F)([\mathbf{X}_u]) \right) v_+ \left(\frac{K(\mathbf{X}_u)}{\varepsilon^r} \right) du \right| \right] = 0. \quad (4.77)$$

Finally, combine (4.67), (4.68), and (4.77) to obtain (4.65), completing the proof of Proposition 4.7. $\square$

Remark 4.6. Notice that we require

$$2q < r < \min \left\{ p, \frac{1-5q}{n+1} \right\}.$$

Therefore, we need to make sure that $q < \min \left\{ \frac{p}{2}, \frac{1}{2n+7} \right\}$. Recall that $p = \frac{1}{n+1}$, which implies that we need $q < \frac{1}{2n+7}$.

Proposition 4.8. Fix $0 \leq t_1 < t_2 < \dots < t_m \leq s < t$ and $h_1, \dots, h_m \in C_b(\mathbb{R}^2)$. If $F \in \mathcal{D}^*$, then

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} (\mathbb{M}G_i^\varepsilon)(\mathbf{X}_s) ds \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] = 0, \quad i = 0, 1, 2. \quad (4.78)$$

Proof. Notice that

$$\begin{aligned} (\mathbb{M}G_0^\varepsilon)(\mathbf{x}) &= \frac{1}{\varepsilon^{p+1}} v'_+ \\ &\quad \times \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x}) \right) (\mathbb{M}(\nabla K, b))(\mathbf{x}) = 0, \\ (\mathbb{M}G_1^\varepsilon)(\mathbf{x}) &= \frac{1}{\varepsilon^p} v'_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \\ &\quad \times \left((\mathbb{M}\langle dF_{\mathbf{V}}^\varepsilon, dK \rangle)(\mathbf{x}) - (F_K^\varepsilon)'(K(\mathbf{x})) (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}) \right), \\ (\mathbb{M}G_2^\varepsilon)(\mathbf{x}) &= \frac{1}{2\varepsilon^{2p}} v''_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x}) \right) (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x}). \end{aligned} \quad (4.79)$$

For $\varepsilon > 0$ we define two functions $f_1^\varepsilon, f_2^\varepsilon \in C^4(\bar{\mathbf{I}} \cup \{\mathbf{x} : \varepsilon^p < K(\mathbf{x}) < 2\varepsilon^p\})$ by

$$f_1^\varepsilon(\mathbf{x}) := (\mathbb{M}\langle dF_{\mathbf{V}}^\varepsilon, dK \rangle)(\mathbf{x}) - (F_K^\varepsilon)'(K(\mathbf{x})) (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x})$$

and

$$f_2^\varepsilon(\mathbf{x}) := \frac{F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})}{K(\mathbf{x})} (\mathbb{M}(dK, dK))(\mathbf{x}).$$

We also define

$$\omega_1(h) := \frac{v'_+(h)}{h^n}, \quad \omega_2(h) := \frac{v''_+(h)}{2h^{n-1}}, \quad h \in \mathbb{R}.$$

Then, for all $\mathbf{x} \in \bar{\mathbf{I}}$,

$$\begin{aligned} \frac{1}{\varepsilon^p} v'_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) &= \varepsilon^{-p(n+1)} \omega_1 \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) (K(\mathbf{x}))^n \\ &= \varepsilon^{-1} \omega_1 \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) d_n(K(\mathbf{x})) \end{aligned}$$

and

$$\begin{aligned} \frac{1}{2\varepsilon^{2p}} K(x) v''_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) &= \varepsilon^{-2p} K(\mathbf{x}) \omega_2 \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right)^{n-1} \\ &= \varepsilon^{-1} \omega_2 \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) d_n(K(\mathbf{x})). \end{aligned}$$

Therefore,

$$(\mathbb{M}G_1^\varepsilon)(\mathbf{x}) = \varepsilon^{-1} \omega_1 \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) d_n(K(\mathbf{x})) f_1^\varepsilon(\mathbf{x})$$

and

$$(\mathbb{M}G_2^\varepsilon)(\mathbf{x}) = \varepsilon^{-1} \omega_2 \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) d_n(K(\mathbf{x})) f_2^\varepsilon(\mathbf{x}).$$

Notice that

$$\begin{aligned} (\mathbb{M}G_i^\varepsilon)(\mathbf{x}) &= \varepsilon^{-1} \left(f_i^\varepsilon(\mathbf{x}) - (\mathbb{A}_0 f_i^\varepsilon)([\mathbf{x}]) \right) d_n(K(\mathbf{x})) \omega_i \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \\ &\quad + \varepsilon^{-1} (\mathbb{A}_0 f_i^\varepsilon)([\mathbf{x}]) d_n(K(\mathbf{x})) \omega_i \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right), \quad i = 1, 2. \end{aligned} \quad (4.80)$$

Define

$$\gamma_i(\varepsilon) := \sup_{\varepsilon^p < h < 2\varepsilon^p} \left| (\mathbb{A}_0 f_i^\varepsilon)([K^{-1}(h)]) \right| = \sup_{\varepsilon^p < h < 2\varepsilon^p} \left| (\mathbb{A}_K f_i^\varepsilon)(h) \right|, \quad i = 1, 2.$$

Notice that there exists a positive constant C^* such that

$$\left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} (\mathbb{A}_0 f_i^\varepsilon)([\mathbf{X}_u]) d_n(K(\mathbf{X}_u)) \omega_i \left(\frac{K(\mathbf{X}_u)}{\varepsilon^p} \right) du \right] \right| < C^* \varepsilon \gamma_i(\varepsilon). \quad (4.81)$$

Now combine (4.80), (4.81) and Proposition 3.3 with ω_i for ω and with $f_i^\varepsilon, i = 1, 2$, for f to see that

$$\begin{aligned} \left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} (\mathbb{M}G_i^\varepsilon)(\mathbf{X}_s) ds \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right| \\ < C(1+t)\varepsilon^p \|f_i^\varepsilon\|_{C^5(\mathcal{E})} + C^* \gamma_i(\varepsilon). \end{aligned} \quad (4.82)$$

N. V. O'Bryant

Lemma 4.12.

$$\sup_{\varepsilon > 0} \left(\varepsilon^{6q} \|f_1^\varepsilon\|_{C^5(\mathcal{E})} \right) < \infty \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \gamma_1(\varepsilon) = 0.$$

Proof. Recall that

$$f_1^\varepsilon(\mathbf{x}) := (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle)(\mathbf{x}) - (F_K^\varepsilon)'(K(\mathbf{x})) (\mathbb{M} \langle dK, dK \rangle)(\mathbf{x})$$

and

$$\left(A_K f_1^\varepsilon \right)(h) = \left(A_K (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle) \right)(h) - (F_K^\varepsilon)'(h) \left(A_K (\mathbb{M} \langle dK, dK \rangle) \right)(h).$$

Notice that all derivatives of f_1^ε of order ≤ 5 involve derivatives of $F_{\mathbf{V}}^\varepsilon$ and F_K^ε of order ≤ 6 . Therefore, by Lemma 4.9 and Lemma 4.10, there exists a constant $C > 0$ such that

$$\|f_1^\varepsilon\|_{C^5(\mathcal{E})} < C\varepsilon^{-6q}.$$

Using the gluing requirement $\underline{G}F = \overline{G}F$ on the function $F \in \mathcal{D}^*$, where

$$\underline{G}F = \left(A^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right)(0)$$

and

$$\overline{G}F = F_K'(0) \left(A_K (\mathbb{M} \langle dK, dK \rangle) \right)(0),$$

we see that

$$\begin{aligned} \left| \left(A_K f_1^\varepsilon \right)(h) \right| &\leq \left| \left(A^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle) \right)(h) - \left(A^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle) \right)(0) \right| \\ &\quad + \left| \left(A^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle) \right)(0) - \left(A^{\text{Tr}} (\mathbb{M} \langle dF_{\mathbf{V}}, dK \rangle) \right)(0) \right| \\ &\quad + \left| F_K'(0) \left(A_K (\mathbb{M} \langle dK, dK \rangle) \right)(0) - (F_K^\varepsilon)'(0) \left(A_K (\mathbb{M} \langle dK, dK \rangle) \right)(0) \right| \\ &\quad + \left| (F_K^\varepsilon)'(0) \left(A_K (\mathbb{M} \langle dK, dK \rangle) \right)(0) - (F_K^\varepsilon)'(h) \left(A_K (\mathbb{M} \langle dK, dK \rangle) \right)(h) \right|. \end{aligned}$$

for all $h \in (-a, a)$. Using properties of the trace operators, Lemma 4.11, and the approximation results of Lemma 4.9 and Lemma 4.10, we can show that there exists a constant $C > 0$ such that

$$|\gamma_1(\varepsilon)| \leq C \left(\|F_{\mathbf{V}}^\varepsilon\|_{\mathbb{H}^2(\mathbf{V})} \sqrt{|h|} + \|F_{\mathbf{V}}^\varepsilon - F_{\mathbf{V}}\|_{\mathbb{H}^2(\mathbf{V})} + \|F_K^\varepsilon - F_K\|_{C^1((0, K^*))} + \varepsilon^{p-2q} \right)$$

for all $h \in (-a, a)$. With $q < p/2$, the inequality above implies that $\lim_{\varepsilon \rightarrow 0} \gamma_1(\varepsilon) = 0$.

□

Before finishing the proof of Proposition 4.8, we will need several auxiliary results.

Lemma 4.13. *Let $f \in C^n(\mathbb{R})$ for some integer $n \geq 2$.*

There exists a function $R_{f,\mathbf{U}} \in C^{n-2}(\mathbb{R})$ such that

$$f(h) = f(0) + hf'(h) + h^2 R_{f,\mathbf{U}}(h), \quad (4.83)$$

For each integer m , $0 < m \leq n$, there exists a constant $C_m > 0$ such that

$$\|R_{f,\mathbf{U}}\|_{C^{m-2}([-a,a])} \leq C_m \|f\|_{C^m([-a,a])}. \quad (4.84)$$

A very simple proof of Lemma 4.13 can be found in [?] (Lemma 9.4).

Lemma 4.14. *Let $f \in C^n(\mathbb{R}^2)$ for some integer $n \geq 2$, and suppose that $f(\mathbf{x}) \equiv f_0$ for $\mathbf{x} \in \partial\mathbf{V}$. Then*

$$(\mathbb{M}\langle df, dK \rangle)(\mathbf{x}) = \frac{(\nabla f, \nabla K)(\mathbf{x})}{\|\nabla K(\mathbf{x})\|^2} (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x})$$

for all $\mathbf{x} \in \partial\mathbf{V}$. There exists a function $R_{f,\mathbf{V}} \in C^{n-2}(\mathbb{R}^2)$ such that for every $\mathbf{x} \in \mathcal{E}$,

$$f(\mathbf{x}) = f_0 + K(\mathbf{x}) \frac{(\mathbb{M}\langle df, dK \rangle)(\mathbf{x})}{(\mathbb{M}\langle dK, dK \rangle)(\mathbf{x})} + K^2(\mathbf{x}) R_{f,\mathbf{V}}(\mathbf{x}). \quad (4.85)$$

For each integer m , $0 < m \leq n$, there exists a constant $C_m > 0$ such that

$$\|R_{f,\mathbf{V}}\|_{C^{m-2}(\overline{\mathcal{E}})} \leq C_m \|f\|_{C^m(\overline{\mathcal{E}})}. \quad (4.86)$$

Proof. Let $\mathbf{x} \in \partial\mathbf{V}$, then $(\nabla f, \nabla^\perp K)(\mathbf{x}) = 0$. By projecting ∇f onto two orthogonal directions of ∇K and $\nabla^\perp K$, we see that

$$\begin{aligned} \nabla f(\mathbf{x}) &= \frac{(\nabla f, \nabla K)(\mathbf{x})}{\|\nabla K(\mathbf{x})\|^2} \nabla K(\mathbf{x}) + \frac{(\nabla f, \nabla^\perp K)(\mathbf{x})}{\|\nabla^\perp K(\mathbf{x})\|^2} \nabla^\perp K(\mathbf{x}) \\ &= \frac{(\nabla f, \nabla K)(\mathbf{x})}{\|\nabla K(\mathbf{x})\|^2} \nabla K(\mathbf{x}). \end{aligned}$$

Therefore, using the definition of $\langle df, dg \rangle_t$, we see that

$$(\mathbb{M}\langle df, dK \rangle)(\mathbf{x}) = \frac{(\nabla f, \nabla K)(\mathbf{x})}{\|\nabla K(\mathbf{x})\|^2} (\mathbb{M}\langle dK, dK \rangle)(\mathbf{x})$$

for all $\mathbf{x} \in \partial\mathbf{V}$.

Using the flow ζ_t on $\mathcal{E}$ introduced in (3.37) we consider the function $F : t \mapsto f(\zeta_{-tK(\mathbf{x})}(\mathbf{x}))$. Notice that

$$\dot{\zeta}_{-tK(\mathbf{x})}(\mathbf{x}) = -K(\mathbf{x}) \frac{\nabla K}{\|\nabla K\|^2}(\zeta_{-tK(\mathbf{x})}(\mathbf{x})),$$

$F(0) = f(\zeta_0(\mathbf{x})) = f(\mathbf{x})$, and $F(1) = f(\zeta_{-K(\mathbf{x})}(\mathbf{x})) = f_0$. Using the identity

$$F(t) = F(s) + F'(t)(t-s) + \int_s^t (s-r) F''(r) dr$$

N. V. O'Bryant

with $t = 0$ and $s = 1$, we obtain

$$\begin{aligned} f(\mathbf{x}) &= f_0 + K(\mathbf{x}) \frac{(\nabla f, \nabla K)(\mathbf{x})}{\|\nabla K(\mathbf{x})\|^2} + K^2(\mathbf{x}) R_{f, \mathbf{V}}(\mathbf{x}) \\ &= f_0 + K(\mathbf{x}) \frac{(\mathbb{M}\langle df, dK \rangle)(\mathbf{x})}{(\mathbb{M}\langle dK, dK \rangle)(\mathbf{x})} + K^2(\mathbf{x}) R_{f, \mathbf{V}}(\mathbf{x}), \end{aligned}$$

for all $\mathbf{x} \in \mathcal{E}$, where

$$R_{f, \mathbf{V}}(\mathbf{x}) = \int_0^1 (t-1) \left(\nabla \left(\frac{(\nabla f, \nabla K)(\mathbf{x})}{\|\nabla K(\mathbf{x})\|^2} \right), \frac{\nabla K}{\|\nabla K\|^2} \right) (\zeta_{-tK(\mathbf{x})}(\mathbf{x})) dt.$$

Let $0 < m \leq n$, then from this expression for $R_{f, \mathbf{V}}(\mathbf{x})$ we see that there exists a constant $C_m > 0$ such that the bound (4.86) holds. $\square$

Lemma 4.15. *There exists a function $f_3^\varepsilon \in C^5(\mathcal{E})$ satisfying $\mathbb{A}_0 f_3^\varepsilon \equiv 0$, and*

$$\sup_{\varepsilon > 0} \varepsilon^{8q} \|f_3^\varepsilon\|_{C^5(\mathcal{E})} < \infty, \quad (4.87)$$

such that for all $\mathbf{x} \in \mathcal{E}$,

$$(\nabla F_{\mathbf{V}}^\varepsilon(\mathbf{x}), \nabla^\perp K(\mathbf{x})) = K(\mathbf{x}) f_3^\varepsilon(\mathbf{x}). \quad (4.88)$$

Proof. By Lemma 4.14,

$$F_{\mathbf{V}}^\varepsilon(\mathbf{x}) = F(\partial \mathbf{V}) + K(\mathbf{x}) \frac{(\mathbb{M}\langle dF_{\mathbf{V}}^\varepsilon, dK \rangle)(\mathbf{x})}{(\mathbb{M}\langle dK, dK \rangle)(\mathbf{x})} + K^2(\mathbf{x}) R_{F_{\mathbf{V}}^\varepsilon, \mathbf{V}}(\mathbf{x})$$

for all $\mathbf{x} \in \mathcal{E}$, which implies that

$$\begin{aligned} (\nabla F_{\mathbf{V}}^\varepsilon(\mathbf{x}), \nabla^\perp K(\mathbf{x})) &= K(\mathbf{x}) \left(\nabla \left(\frac{(\mathbb{M}\langle dF_{\mathbf{V}}^\varepsilon, dK \rangle)(\mathbf{x})}{(\mathbb{M}\langle dK, dK \rangle)(\mathbf{x})} \right), \nabla^\perp K(\mathbf{x}) \right) \\ &\quad + K^2(\mathbf{x}) (\nabla R_{F_{\mathbf{V}}^\varepsilon, \mathbf{V}}(\mathbf{x}), \nabla^\perp K(\mathbf{x})). \end{aligned}$$

Take

$$f_3^\varepsilon(\mathbf{x}) = \left(\nabla^\perp K(\mathbf{x}), \nabla \left(\frac{(\mathbb{M}\langle dF_{\mathbf{V}}^\varepsilon, dK \rangle)(\mathbf{x})}{(\mathbb{M}\langle dK, dK \rangle)(\mathbf{x})} \right) \right) + K(\mathbf{x}) (\nabla^\perp K(\mathbf{x}), \nabla R_{F_{\mathbf{V}}^\varepsilon, \mathbf{V}}(\mathbf{x})).$$

Then (4.88) holds. From this identity and from (4.89) we see that for any $\mathbf{x} \in \mathcal{E}$

$$(\mathbb{A}_0 (K(\mathbf{x}) f_3^\varepsilon))([\mathbf{x}]) = K(\mathbf{x}) (\mathbb{A}_0 f_3^\varepsilon)([\mathbf{x}]) \equiv 0.$$

Therefore, $\mathbb{A}_0 f_3^\varepsilon \equiv 0$.

To verify that (4.87) holds, we use inequality (4.86) and Lemma 4.9. This finishes the proof of Lemma 4.15. $\square$

Lemma 4.16.

$$\sup_{\varepsilon > 0} \left(\varepsilon^{7q} \|f_2^\varepsilon\|_{C^5(\mathcal{E})} \right) < \infty \quad \text{and} \quad \lim_{\varepsilon \rightarrow 0} \gamma_2(\varepsilon) = 0.$$

Proof. Recall that

$$f_2^\varepsilon(\mathbf{x}) := \frac{F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})}{K(\mathbf{x})} (\mathbb{M} \langle dK, dK \rangle)(\mathbf{x}),$$

$F_K^\varepsilon(0) = F(\Gamma_{\partial \mathbf{V}})$, and $F_{\mathbf{V}}^\varepsilon(\mathbf{x}) = F(\Gamma_{\partial \mathbf{V}})$ for $\mathbf{x} \in \partial \mathbf{V}$.

By Lemma 4.13 and Lemma 4.14, if $\mathbf{x} \in \mathcal{E}$ then

$$\begin{aligned} \frac{F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})}{K(\mathbf{x})} &= \frac{F_K^\varepsilon(K(\mathbf{x})) - F(\Gamma_{\partial \mathbf{V}})}{K(\mathbf{x})} - \frac{F_{\mathbf{V}}^\varepsilon(\mathbf{x}) - F(\Gamma_{\partial \mathbf{V}})}{K(\mathbf{x})} \\ &= (F_K^\varepsilon)'(K(\mathbf{x})) + K(\mathbf{x}) R_{F_K^\varepsilon, \mathbf{U}}(K(\mathbf{x})) \\ &\quad - \frac{(\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle)(\mathbf{x})}{(\mathbb{M} \langle dK, dK \rangle)(\mathbf{x})} - K(\mathbf{x}) R_{F_{\mathbf{V}}^\varepsilon, \mathbf{V}}(\mathbf{x}) \end{aligned}$$

Now, by Lemma 4.10, Lemma 4.9, and applying inequalities (4.84) and (4.86), there exists a constant $C > 0$ such that

$$\|f_2^\varepsilon\|_{C^3(\mathcal{E})} < C\varepsilon^{-7q}.$$

Similarly, notice that

$$\begin{aligned} f_2^\varepsilon(\mathbf{x}) - f_1^\varepsilon(\mathbf{x}) &= \left(\frac{F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})}{K(\mathbf{x})} - (F_K^\varepsilon)'(K(\mathbf{x})) \right) (\mathbb{M} \langle dK, dK \rangle)(\mathbf{x}) \\ &\quad - (\mathbb{M} \langle dF_{\mathbf{V}}^\varepsilon, dK \rangle)(\mathbf{x}), \end{aligned}$$

and so there exists a constant $C > 0$ such that

$$|f_2^\varepsilon(\mathbf{x}) - f_1^\varepsilon(\mathbf{x})| \leq C\varepsilon^{p-2q}$$

for all $\mathbf{x} \in \bar{\mathbf{I}} \cap \{\mathbf{y} : \varepsilon^p < K(\mathbf{y}) < 2\varepsilon^p\}$. Now,

$$\gamma_2(\varepsilon) \leq \gamma_1(\varepsilon) + C\varepsilon^{p-2q}.$$

Therefore, by Lemma 4.12, and since $q < p/2$, $\lim_{\varepsilon \rightarrow 0} \gamma_2(\varepsilon) = 0$. $\square$

From Lemma 4.12 and Lemma 4.16, $\lim_{\varepsilon \rightarrow 0} \gamma_1(\varepsilon) = \lim_{\varepsilon \rightarrow 0} \gamma_2(\varepsilon) = 0$. Therefore, in the case of $i = 1, 2$, (4.78) follows from (4.82). The case $i = 0$ is trivial due to the fact that $(\mathbb{M} G_0^\varepsilon)(\mathbf{x}) \equiv 0$. This completes the proof of Proposition 4.8. $\square$

Proposition 4.9. Fix $0 \leq t_1 < t_2 < \dots < t_m \leq s < t$ and $h_1, \dots, h_m \in C_b(\mathbb{R}^2)$. If $F \in \mathcal{D}^*$ then

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} (\mathbb{M} G_3^\varepsilon)(\mathbf{X}_s) ds \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] = 0.$$

Proof. Notice that

$$\begin{aligned} (\mathbb{M} G_3^\varepsilon)(\mathbf{x}) &= \frac{1}{\varepsilon} \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \right) (\nabla F_{\mathbf{V}}^\varepsilon(\mathbf{x}), \nabla^\perp H(\mathbf{x})) \\ &= \frac{1}{\varepsilon} \left(1 - v_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \right) n d_{n-1}(K(\mathbf{x})) (\nabla F_{\mathbf{V}}^\varepsilon(\mathbf{x}), \nabla^\perp K(\mathbf{x})) \end{aligned}$$

N. V. O'Bryant

and

$$\begin{aligned}
(\mathbb{A}_0 (\nabla F_{\mathbf{V}}^\varepsilon, \nabla^\perp K)) ([\mathbf{x}]) &= \frac{1}{\eta(K(\mathbf{x}))} \int_0^{\eta(K(\mathbf{x}))} (\nabla F_{\mathbf{V}}^\varepsilon, \nabla^\perp K) (\phi_t(\mathbf{x})) dt \\
&= \frac{1}{\eta(K(\mathbf{x}))} \int_0^{\eta(K(\mathbf{x}))} \frac{d}{dt} F_{\mathbf{V}}^\varepsilon(\phi_t(\mathbf{x})) dt \\
&= F_{\mathbf{V}}^\varepsilon(\phi_{\eta(K(\mathbf{x}))}(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x}) = 0.
\end{aligned} \tag{4.89}$$

Let f_3^ε be a function from Lemma 4.15, and define

$$\omega_3(h) := n(1 - v_+(h)), \quad h \in \mathbb{R}.$$

Then,

$$(\mathbb{M}G_3^\varepsilon)(\mathbf{x}) = \varepsilon^{-1} \omega_3 \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \mathbf{d}_n(K(\mathbf{x})) f_3^\varepsilon(\mathbf{x}). \tag{4.90}$$

Now combine (4.90), and Proposition 3.3 with ω_3 for ω and with f_3^ε for f to see that

$$\left| \mathbb{E}^\varepsilon \left[\int_{s \wedge \tau}^{t \wedge \tau} (\mathbb{M}G_3^\varepsilon)(\mathbf{X}_s) ds \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] \right| < C \varepsilon^p (1+t) \|f_3^\varepsilon\|_{C^5(\mathcal{E})}. \tag{4.91}$$

Thus, Proposition 4.9 holds. $\square$

Proposition 4.10. *Fix $F \in \mathcal{D}^*$ and $t > 0$. Then*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\left| \int_0^{t \wedge \tau} (\mathbb{M}R^\varepsilon)(\mathbf{X}_s) ds \right| \right] = 0.$$

Proof. Consider

$$(\mathbb{M}R^\varepsilon)(\mathbf{x}) = \frac{1}{\varepsilon^p} v'_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \left(F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x}) \right) (\mathbb{M}(\mathcal{L}K))(\mathbf{x})$$

Define

$$\beta_K(\varepsilon) := \sup_{\substack{\mathbf{x} \in \bar{\mathbf{I}} \\ \varepsilon^p \leq K(\mathbf{x}) \leq 2\varepsilon^p}} |F_K^\varepsilon(K(\mathbf{x})) - F_{\mathbf{V}}^\varepsilon(\mathbf{x})|,$$

which satisfies

$$\beta_K(\varepsilon) \leq \sup_{\substack{\mathbf{x} \in \bar{\mathbf{I}} \\ \varepsilon^p \leq K(\mathbf{x}) \leq 2\varepsilon^p}} \left(|F_K^\varepsilon(K(\mathbf{x})) - F(\mathbf{\Gamma}_{\partial \mathbf{V}})| + |F_{\mathbf{V}}^\varepsilon(\mathbf{x}) - F(\mathbf{\Gamma}_{\partial \mathbf{V}})| \right). \tag{4.92}$$

Notice that

$$F_K^\varepsilon(K(\mathbf{x})) - F(\mathbf{\Gamma}_{\partial \mathbf{V}}) = F_K^\varepsilon(K(\mathbf{x})) - F_K^\varepsilon(0) = (F_K^\varepsilon)'(0)K(\mathbf{x}) + o(|K(\mathbf{x})|).$$

From this expression and Lemma 4.10, there exist constants $C_1, C_{11} > 0$ such that

$$|F_K^\varepsilon(K(\mathbf{x})) - F(\mathbf{\Gamma}_{\partial \mathbf{V}})| < C_1 \varepsilon^p \|F_K^\varepsilon\|_{C^1([0, K^*])} < C_{11} \varepsilon^{p-q}. \tag{4.93}$$

By the definition of K , if $\mathbf{x}^* \in \partial\mathbf{V}$, then $K(\mathbf{x}^*) = 0$, and $\nabla K(\mathbf{x}^*) \neq 0$. Therefore, there exists $\mathbf{x}^* \in \partial\mathbf{V}$, such that

$$\begin{aligned} F_{\mathbf{V}}^{\varepsilon}(\mathbf{x}) - F(\Gamma_{\partial\mathbf{V}}) &= F_{\mathbf{V}}^{\varepsilon}(\mathbf{x}) - F_{\mathbf{V}}^{\varepsilon}(\mathbf{x}^*) \\ &= (\nabla F_{\mathbf{V}}^{\varepsilon}(\mathbf{x}^*), \mathbf{x} - \mathbf{x}^*) + o(\|\mathbf{x} - \mathbf{x}^*\|) \\ &= \|\nabla F_{\mathbf{V}}^{\varepsilon}(\mathbf{x}^*)\| \|\mathbf{x} - \mathbf{x}^*\| + o(\|\mathbf{x} - \mathbf{x}^*\|) \\ &= \frac{\|\nabla F_{\mathbf{V}}^{\varepsilon}(\mathbf{x}^*)\|}{\|\nabla K(\mathbf{x}^*)\|} K(\mathbf{x}) + o(|K(\mathbf{x})|). \end{aligned}$$

From this expression and Lemma 4.9, there exist constants $C_2, C_{22} > 0$ such that

$$|F_{\mathbf{V}}^{\varepsilon}(\mathbf{x}) - F(\Gamma_{\partial\mathbf{V}})| < C_2 \varepsilon^p \|F_{\mathbf{V}}^{\varepsilon}\|_{C^1(\bar{\mathbf{I}})} < C_{22} \varepsilon^{p-q}. \quad (4.94)$$

Now, combine inequalities (4.92), (4.93), and (4.94) to conclude that $\beta_K(\varepsilon) \leq (C_{11} + C_{22}) \varepsilon^{p-q}$ for all $\varepsilon > 0$. Therefore, since

$$\left| v'_+ \left(\frac{K(\mathbf{x})}{\varepsilon^p} \right) \right| \leq C_w \chi_{[\varepsilon^p, 2\varepsilon^p]}(K(\mathbf{x}))$$

for some $C_w > 0$ and

$$\sup_{\mathbf{x} \in \bar{\mathbf{I}}} |(\mathbb{M}(\mathcal{L}K))(\mathbf{x})| \leq C_K$$

for some $C_K > 0$, there exists a positive constant $C := C_w C_K (C_{11} + C_{22})$ such that

$$\mathbb{E}^{\varepsilon} \left[\left| \int_0^{t \wedge \tau} (\mathbb{M}R^{\varepsilon})(\mathbf{X}_u) du \right| \right] \leq C \varepsilon^{p-q} \mathbb{E}^{\varepsilon} \left[\frac{1}{\varepsilon^p} \int_0^{t \wedge \tau} \chi_{[\varepsilon^p, 2\varepsilon^p]}(K(\mathbf{X}_u)) du \right].$$

This inequality and Lemma 3.5 give the desired result. $\square$

Proposition 4.11. Fix $0 \leq t_1 < t_2 < \dots < t_m \leq s < t$ and $h_1, \dots, h_m \in C_b(\mathbb{R}^2)$. If $F \in \mathcal{D}^*$, then

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E}^{\varepsilon} \left[\int_0^{t \wedge \tau} \left((\mathbb{M}(\mathcal{L}^{\varepsilon} F^{\varepsilon}))(\mathbf{X}_s) - (\mathcal{L}^* F)([\mathbf{X}_s]) \right) ds \prod_{k=1}^m h_k(\mathbf{X}_{t_k}) \right] = 0. \quad (4.95)$$

Proof. Given $\mathbf{x} \in \bar{\mathbf{I}}$, consider

$$\begin{aligned} (\mathbb{M}(\mathcal{L}^{\varepsilon} F^{\varepsilon}))(\mathbf{x}) - (\mathcal{L}^* F)([\mathbf{x}]) &= \left((\mathbb{M}L^{\varepsilon})(\mathbf{x}) - (\mathcal{L}^* F)([\mathbf{x}]) \right) \\ &\quad + (\mathbb{M}K^{\varepsilon})(\mathbf{x}) + \sum_{k=0}^3 (\mathbb{M}G_k^{\varepsilon})(\mathbf{x}) + (\mathbb{M}R^{\varepsilon})(\mathbf{x}). \end{aligned}$$

Now, combine Proposition 4.7, Proposition 4.8, Proposition 4.9, and Proposition 4.10 to verify that (4.95) holds. $\square$

Finally, we can prove the desired statement about the limiting martingale problem.

N. V. O'Bryant

Theorem 4.1. Fix $\mathbf{x} \in \bar{\mathbf{I}}$. Let $\mathbb{P}^* \in \mathcal{P}(C([0, \infty); \mathbf{\Gamma}))$ be a cluster point of the $\mathbb{P}^{\varepsilon, *}_s$. Then $\mathbb{P}^*$ satisfies the martingale problem for $(\mathcal{L}^*, \delta_{[\mathbf{x}]})$.

Proof. Let $\mathbb{E}^\varepsilon$ be the expectation operator associated with the law $\mathbb{P}^\varepsilon$ and let $\mathbb{E}^*$ be the expectation operator associated with the law $\mathbb{P}^*$.

We know that every $\mathbb{P}^\varepsilon \in \mathcal{P}(C[0, \infty); \mathbb{R}^2)$ is a solution of the stopped martingale problem for $(\mathcal{L}_{t/\varepsilon^2}^\varepsilon, \delta_{\mathbf{x}}, \mathbf{I})$, and that $F^\varepsilon \in C^2(\mathbb{R}^2) \subset \mathcal{D}^\varepsilon$. Therefore, we know that for any $0 \leq s < t$, and $\varepsilon > 0$,

$$F^\varepsilon(\mathbf{X}_{t \wedge \tau}) - F^\varepsilon(\mathbf{X}_{s \wedge \tau}) - \int_{s \wedge \tau}^{t \wedge \tau} (\mathcal{L}_{u/\varepsilon^2}^\varepsilon F^\varepsilon)(\mathbf{X}_u) du \quad (4.96)$$

is a $\mathbb{P}^\varepsilon$ -martingale. We also know that $\mathbb{P}^\varepsilon\{\mathbf{X}_0 = \mathbf{x}\} = 1$.

Our theorem claims that $\mathbb{P}^*\{\mathbf{Y}_0 = [\mathbf{x}]\} = 1$ and if $F \in \mathcal{D}^*$, $0 \leq t_1 < t_2 < \dots < t_m \leq s < t$, and $h_1^*, \dots, h_m^* \in C_b(\mathbf{\Gamma})$ then

$$\mathbb{E}^* \left[\left\{ F(\mathbf{Y}_t) - F(\mathbf{Y}_s) - \int_s^t (\mathcal{L}^* F)(\mathbf{Y}_u) du \right\} \prod_{k=1}^m h_k^*(\mathbf{Y}_{t_k}) \right] = 0. \quad (4.97)$$

Notice that if $B \subset \mathbf{\Gamma}$ is any set which does not contain $[\mathbf{x}]$, then $\pi^{-1}(B) \subset \bar{\mathbf{I}}$ does not contain $\mathbf{x}$, and

$$\mathbb{P}^*\{\mathbf{Y}_0 \in B\} \leq \overline{\lim}_{\varepsilon \rightarrow 0} \mathbb{P}^\varepsilon\{\mathbf{Y}_0 \in B\} = \overline{\lim}_{\varepsilon \rightarrow 0} \mathbb{P}^\varepsilon\{\mathbf{X}_0 \in \pi^{-1}(B)\} = 0.$$

Therefore $\mathbb{P}^*\{\mathbf{Y}_0 = [\mathbf{x}]\} = 1$.

By the definition of $\mathcal{D}^*$ we know that $(\mathcal{L}^* F)(\mathbf{\Gamma}_{\partial \mathbf{I}}) = 0$, which implies that for any $0 \leq s < t$,

$$\int_{s \wedge \tau}^{t \wedge \tau} (\mathcal{L}^* F)(\mathbf{Y}_s) ds = \int_s^t (\mathcal{L}^* F)(\mathbf{Y}_s) ds.$$

Now,

$$\begin{aligned} & \mathbb{E}^* \left[\left\{ F(\mathbf{Y}_t) - F(\mathbf{Y}_s) - \int_s^t (\mathcal{L}^* F)(\mathbf{Y}_u) du \right\} \prod_{k=1}^m h_k^*(\mathbf{Y}_{t_k}) \right] \\ &= \lim_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\left\{ F(\mathbf{Y}_t) - F(\mathbf{Y}_s) - \int_s^t (\mathcal{L}^* F)(\mathbf{Y}_u) du \right\} \prod_{k=1}^m h_k^*(\mathbf{Y}_{t_k}) \right] \\ &= \lim_{\varepsilon \rightarrow 0} \mathbb{E}^\varepsilon \left[\left\{ F \circ \pi(\mathbf{X}_{t \wedge \tau}) - F \circ \pi(\mathbf{X}_{s \wedge \tau}) - \int_{s \wedge \tau}^{t \wedge \tau} (\mathcal{L}^* F)(\mathbf{Y}_u) du \right\} \prod_{k=1}^m h_k^*(\mathbf{Y}_{t_k}) \right]. \end{aligned} \quad (4.98)$$

Notice that

$$\begin{aligned}
& F \circ \pi(\mathbf{X}_{t \wedge \tau}) - F \circ \pi(\mathbf{X}_{s \wedge \tau}) - \int_{s \wedge \tau}^{t \wedge \tau} (\mathcal{L}^* F)(\mathbf{Y}_u) du \\
&= \left(F \circ \pi(\mathbf{X}_{t \wedge \tau}) - F^\varepsilon(\mathbf{X}_{t \wedge \tau}) \right) - \left(F \circ \pi(\mathbf{X}_{s \wedge \tau}) - F^\varepsilon(\mathbf{X}_{s \wedge \tau}) \right) \\
&+ F^\varepsilon(\mathbf{X}_{t \wedge \tau}) - F^\varepsilon(\mathbf{X}_{s \wedge \tau}) - \int_{s \wedge \tau}^{t \wedge \tau} \mathcal{L}_{u/\varepsilon^2}^\varepsilon F^\varepsilon(\mathbf{X}_u) du \\
&+ \int_{s \wedge \tau}^{t \wedge \tau} \left(\left(\mathcal{L}_{u/\varepsilon^2}^\varepsilon F^\varepsilon \right)(\mathbf{X}_u) - (\mathbb{M}(\mathcal{L}^\varepsilon F^\varepsilon))(\mathbf{X}_u) \right) du \\
&+ \int_{s \wedge \tau}^{t \wedge \tau} \left((\mathbb{M}(\mathcal{L}^\varepsilon F^\varepsilon))(\mathbf{X}_u) - (\mathcal{L}^* F)(\mathbf{Y}_u) \right) du
\end{aligned}$$

Keeping this expression in mind, we use the statement of the martingale problem (4.96), Lemma 3.1 with $f(\mathbf{x}, t) = \mathcal{L}_t^\varepsilon F^\varepsilon(\mathbf{x})$ concerning the first-level averaging, and Eq. (4.95) to see that (4.98) implies (4.97), and this completes the proof. $\square$

4.5. Uniqueness

We already know that every limit point of the $\mathbb{P}^\varepsilon$ -laws of $\{\mathbf{Y}_t, t \geq 0\}$ satisfies the martingale problem for $(\mathcal{L}^*, \delta_{[\mathbf{x}]})$. Our next step is to verify that the $\mathbb{P}^\varepsilon$ -laws have only one limit point.

Proposition 4.12. *The operator $\mathcal{L}^*$ generates a strongly continuous contraction semigroup on $C(\Gamma)$.*

Proof. To prove this proposition using the Hille-Yosida theorem, we need to show that $\mathcal{D}^*$ is dense in $C(\Gamma)$, that $\mathcal{L}^*$ satisfies the positive maximum principle, and that the range of $\lambda I - \mathcal{L}^*$ is dense in $C(\Gamma)$ for some $\lambda > 0$.

Lemma 4.17. *$\mathcal{D}^*$ is dense in $C(\Gamma)$.*

Proof. Let $F \in C(\Gamma)$. To prove the statement of this lemma, we will approximate this function with functions $F_n \in \mathcal{D}^*$ in the $\|\cdot\|_{C(\Gamma)}$ -norm. We will construct the F_n 's in two steps.

Given $[\mathbf{x}] \in \Gamma$, take

$$\begin{aligned}
F_n^0([\mathbf{x}]) &:= F([\mathbf{x}]) + \left(F(\Gamma_{\partial \mathbf{V}}) - F([\mathbf{x}]) \right) v(2nK(\mathbf{x})) \\
&+ \left(F(\Gamma_{\partial \mathbf{I}}) - F([\mathbf{x}]) \right) v(2n(K(\mathbf{x}) - K^*)).
\end{aligned}$$

Then,

$$\lim_{n \rightarrow 0} \|F - F_n^0\|_{C(\Gamma)} = 0. \tag{4.99}$$

N. V. O'Bryant

Second, for every n , using the standard mollification, there exists a function $F_n^1 \in C^\infty(\mathbf{V} \cup \mathbf{U})$ such that

$$\sup_{\substack{|K(\mathbf{x})| \geq \frac{1}{2n} \\ |K(\mathbf{x}) - K^*| \geq \frac{1}{2n}}} |F([\mathbf{x}]) - F_n^1([\mathbf{x}])| \leq \frac{1}{n}.$$

Now choose

$$\begin{aligned} F_n([\mathbf{x}]) &:= F_n^1([\mathbf{x}]) + \left(F(\Gamma_{\partial \mathbf{V}}) - F_n^1([\mathbf{x}]) \right) v(2nK(\mathbf{x})) \\ &\quad + \left(F(\Gamma_{\partial \mathbf{U}}) - F_n^1([\mathbf{x}]) \right) v(2n(K(\mathbf{x}) - K^*)). \end{aligned}$$

Then,

$$\lim_{n \rightarrow 0} \|F_n^0 - F_n\|_{C(\Gamma)} = 0. \quad (4.100)$$

Each of the F_n s is a smooth function on $\mathbf{V} \cup \mathbf{U}$ and is a constant in a small neighborhood of $\partial \mathbf{V}$ and in a small neighborhood of $\partial \mathbf{I}$. Thus, $F_n \in \mathcal{D}^*$. Finally, (4.99) and (4.100) imply that

$$\lim_{n \rightarrow 0} \|F - F_n\|_{C(\Gamma)} = 0.$$

Therefore, $\mathcal{D}^*$ is dense in $C(\Gamma)$. $\square$

Lemma 4.18. $\mathcal{L}^*$ satisfies the positive maximum principle.

Proof. By the definition of the positive maximum principle we need to show that whenever $F \in \mathcal{D}^*$, $[\mathbf{x}_0] \in \Gamma$, and $\sup_{[\mathbf{x}] \in \Gamma} F([\mathbf{x}]) = F([\mathbf{x}_0]) \geq 0$, we have $\mathcal{L}^* F([\mathbf{x}_0]) \leq 0$.

Consider $F \in \mathcal{D}^*$, and let it attain its supremum at a point $[\mathbf{x}_0] \in \Gamma$. If $[\mathbf{x}_0] = \Gamma_{\partial \mathbf{I}}$ then, by definition of $\mathcal{D}^*$, we have $\mathcal{L}^* F([\mathbf{x}_0]) = 0$.

Assume that $[\mathbf{x}_0] \in \Gamma_{\mathbf{V}} \cup \Gamma_{\mathbf{U}}$. Recall that the operator $\mathcal{L}^*$ acts locally as a strongly elliptic operator on both $\mathbf{V}$ and $\mathbf{U}$. Therefore, $\mathcal{L}^* F([\mathbf{x}_0]) \leq 0$.

Assume that F attains its supremum at $[\mathbf{x}_0] = \Gamma_{\partial \mathbf{V}}$, then $\overline{GF} \geq 0$ and $\underline{GF} \leq 0$. Since $F \in \mathcal{D}^*$, we know that $\overline{GF} = \underline{GF} = 0$. This implies that

$$F'_K(0) = \frac{\overline{GF}}{\overline{\sigma}^2(0)} = 0.$$

By Lemma 4.10, $F_K \in C^2([0, K^*])$ and $h = 0$ is its local maximum. Therefore, $F''_K(0) \leq 0$. We notice that in this case

$$(\mathcal{L}^* F)([\mathbf{x}_0]) = \lim_{h \downarrow 0} (\mathbb{M}(\mathcal{L}^K F_K))(h) = \overline{b}(0) F'_K(0) + \frac{1}{2} \overline{\sigma}^2(0) F''_K(0) \leq 0.$$

This finishes the proof of Lemma 4.18. $\square$

Lemma 4.19. The range of $\lambda I - \mathcal{L}^*$ is dense in $C(\Gamma)$ for some $\lambda > 0$.

We choose to omit the proof of this lemma since it is very similar to that of Lemma 8.4 in [?].

Lemmas 4.17, 4.18, and 4.19 above allow us to conclude that uniqueness holds for the solution of the martingale problem for $(\mathcal{L}^*, \delta_{[\mathbf{x}]})$. Therefore, the $\mathbb{P}^\varepsilon$ -laws of $\{\mathbf{Y}_t, t \geq 0\}$ converge to the law of a $\mathbf{\Gamma}$ -valued Markov process. $\square$

References